

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



26 JAN 2005



(43) International Publication Date
5 February 2004 (05.02.2004)

PCT

(10) International Publication Number
WO 2004/011443 A1

(51) International Patent Classification⁷: **C07D 239/48**,
239/69, 401/12, 403/12, 405/12, 409/12, 413/12, 413/14,
417/12, A61K 31/5377, 31/506, A61P 11/00, 29/00, 35/00

(21) International Application Number:
PCT/GB2003/003175

(22) International Filing Date: 23 July 2003 (23.07.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0217431.6 27 July 2002 (27.07.2002) GB

(71) Applicant (for AE, AG, AL, AM, AT, AU, AZ, BA, BB, BE, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CY, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, FR, GB, GD, GE, GH, GM, GR, HR, HU, ID, IE, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, SZ, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW only): **ASTRAZENECA AB** [SE/SE]; Sodertalje, S-151 85 (SE).

(71) Applicant (for MG only): **ASTRAZENECA UK LIMITED** [GB/GB]; 15 Stanhope Gate, London, Greater London W1K 1LN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **EBDEN, Mark, Richard** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, Leicestershire LE11 5RH (GB). **MEGHANI, Premji** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough,

Leicestershire LE11 5RH (GB). **BENNION, Collin** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, Leicestershire LE11 5RH (GB). **COOK, Antony, Ronald** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, Leicestershire LE11 5RH (GB). **BONNERT, Roger, Victor** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, Leicestershire LE11 5RH (GB).

(74) Agent: **ASTRAZENECA**; Global Intellectual Property, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PYRIMIDYL SULPHONE AMIDE DERIVATIVES AS CHEMOKINE RECEPTOR MODULATORS



(I)

(57) Abstract: A compound of formula (I), pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof for the treatment of asthma, allergic rhinitis, COPD, inflammatory bowel disease, irritable bowel syndrome, osteoarthritis, osteoporosis, rheumatoid arthritis, psoriasis or cancer.

PYRIMIDYL SULPHONE AMIDE DERIVATIVES AS CHEMOKINE RECEPTOR MODULATORS

The present invention relates to certain heterocyclic compounds, processes and intermediates used in their preparation, pharmaceutical compositions containing them and
5 their use in therapy.

Chemokines play an important role in immune and inflammatory responses in various diseases and disorders, including asthma and allergic diseases, as well as autoimmune pathologies such as rheumatoid arthritis and atherosclerosis. These small secreted molecules are a growing superfamily of 8-14 kDa proteins characterised by a conserved cysteine motif.
10 At the present time, the chemokine superfamily comprises three groups exhibiting characteristic structural motifs, the C-X-C, C-C and C-X₃-C families. The C-X-C and C-C families have sequence similarity and are distinguished from one another on the basis of a single amino acid insertion between the NH-proximal pair of cysteine residues. The C-X₃-C family is distinguished from the other two families on the basis of having a triple amino acid
15 insertion between the NH-proximal pair of cysteine residues.

The C-X-C chemokines include several potent chemoattractants and activators of neutrophils such as interleukin-8 (IL-8) and neutrophil-activating peptide 2 (NAP-2).


The C-C chemokines include potent chemoattractants of monocytes and lymphocytes but not neutrophils. Examples include human monocyte chemotactic proteins 1-
20 3 (MCP-1, MCP-2 and MCP-3), RANTES (Regulated on Activation, Normal T Expressed and Secreted), eotaxin and the macrophage inflammatory proteins 1 α and 1 β (MIP-1 α and MIP-1 β).

The C-X₃-C chemokine (also known as fractalkine) is a potent chemoattractant and activator of microglia in the central nervous system (CNS) as well as of monocytes, T cells,
25 NK cells and mast cells.

Studies have demonstrated that the actions of the chemokines are mediated by subfamilies of G protein-coupled receptors, among which are the receptors designated CCR1, CCR2, CCR2A, CCR2B, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CCR10 and CCR11 (for the C-C family); CXCR1, CXCR2, CXCR3, CXCR4 and CXCR5 (for the C-X-C
30 family) and CX₃CR1 for the C-X₃-C family. These receptors represent good targets for drug development since agents which modulate these receptors would be useful in the treatment of disorders and diseases such as those mentioned above.

-2-

The present invention provides compounds of formula (1), a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof:



5

wherein R¹ is a group selected from C₃₋₇carbocyclyl, C₁₋₈alkyl, C₂₋₆alkenyl and C₂₋₆alkynyl; wherein the group is optionally substituted by 1, 2 or 3 substituents independently selected from fluoro, nitrile, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰,

10 -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, phenyl or heteroaryl; wherein phenyl and heteroaryl are optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, C₁₋₆alkyl and trifluoromethyl;

15 wherein R² is C₃₋₇carbocyclyl, optionally substituted by 1, 2 or 3 substituents independently selected from:

(a) fluoro, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹;

(b) a 3-8 membered ring optionally containing 1, 2 or 3 atoms selected from O, S, -NR⁸

20 and whereby the ring is optionally substituted by C₁₋₃alkyl or fluoro; or

(c) phenyl or heteroaryl, each of which is optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -NR⁸COR⁹, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, C₁₋₆alkyl and trifluoromethyl;

25 or R² is a group selected from C₁₋₈alkyl, C₂₋₆alkenyl or C₂₋₆alkynyl wherein the group is substituted by 1, 2 or 3 substituents independently selected from hydroxy, amino, C₁₋₆alkoxy, C₁₋₆alkylamino, di(C₁₋₆alkyl)amino, *N*-(C₁₋₆alkyl)-*N*-(phenyl)amino, *N*-C₁₋₆alkylcarbonyl,

-3-

N,N-di(C₁₋₆alkyl)carbamoyl, *N*-(C₁₋₆alkyl)-*N*-(phenyl)carbamoyl, carboxy, phenoxycarbonyl, -NR⁸COR⁹, -SO₂R¹⁰, -SO₂NR⁵R⁶ and -NR⁸SO₂R⁹;

wherein R³ is hydrogen or independently R²;

5

R⁴ is hydrogen or a group selected from C₁₋₆alkyl and phenyl, wherein the group is optionally substituted by 1 or 2 substituents independently selected from halo, phenyl, -OR¹¹ and -NR¹²R¹³;

10 R⁵ and R⁶ are independently hydrogen or a group selected from C₁₋₆alkyl and phenyl wherein the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, phenyl, -OR¹⁴, -NR¹⁵R¹⁶, -COOR¹⁴, -CONR¹⁵R¹⁶, -NR¹⁵COR¹⁶, -SO₂R¹⁰, -SONR¹⁵R¹⁶ and NR¹⁵SO₂R¹⁶

or

15 R⁵ and R⁶ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocyclic ring system optionally containing a further heteroatom selected from oxygen and nitrogen atoms, which ring is optionally substituted by 1, 2 or 3 substituents independently selected from phenyl, -OR¹⁴, -COOR¹⁴, -NR¹⁵R¹⁶, -CONR¹⁵R¹⁶, -NR¹⁵COR¹⁶, -SO₂R¹⁰, -SONR¹⁵R¹⁶, NR¹⁵SO₂R¹⁶ or C₁₋₆alkyl (optionally substituted by 1 or 2 substituents independently selected from halo, -NR¹⁵R¹⁶ and -OR¹⁷ groups);

R¹⁰ is hydrogen or a group selected from C₁₋₆alkyl or phenyl, wherein the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, phenyl, -OR¹⁷ and -NR¹⁵R¹⁶; and each of R⁷, R⁸, R⁹, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷ is independently hydrogen,

25 C₁₋₆alkyl or phenyl;

X is hydrogen, halo, cyano, nitro, hydroxy, C₁₋₆alkoxy (optionally substituted by 1 or 2 substituents selected from halo, -OR¹¹ and -NR¹²R¹³), -NR⁵R⁶, -COOR⁷, -NR⁸COR⁹, thio, C₁₋₆alkylthio (optionally substituted by 1 or 2 substituents selected from halo, -OR¹⁷, -NR¹⁵R¹⁶),

30 -SO₂R¹⁰ or a group selected from C₃₋₇carbocyclyl, C₁₋₈alkyl, C₂₋₆alkenyl or C₂₋₆alkynyl, wherein the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶ and -NR⁸SO₂R⁹;

- R^x is trifluoromethyl, -NR⁵R⁶, phenyl, naphthyl, monocyclic or bicyclic heteroaryl wherein a heteroring may be partially or fully saturated and one or more ring carbon atoms may form a carbonyl group, and wherein each phenyl or heteroaryl group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COR⁷, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, C₁₋₆alkyl or trifluoromethyl;
- or R^x is a group selected from C₃₋₇carbocyclyl, C₁₋₈alkyl, C₂₋₆alkenyl and C₂₋₆alkynyl whereby the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COR⁷, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, phenyl or heteroaryl; and wherein each phenyl or heteroaryl group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COR⁷, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, C₁₋₆alkyl or trifluoromethyl;
- or R^x and X together form a 4 to 8-membered sulfonamide ring optionally substituted by 1, 2 or 3 substituents independently selected from halo, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, phenyl or heteroaryl; wherein phenyl and heteroaryl are optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, C₁₋₆alkyl and trifluoromethyl.

Certain compounds of formula (1) are capable of existing in stereoisomeric forms. It will be understood that the invention encompasses all geometric and optical isomers of the compounds of formula (1) and mixtures thereof including racemates.

The synthesis of optically active forms may be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form. Similarly, the above-mentioned activity may be evaluated using the standard laboratory techniques referred to hereinafter.

Within the present invention it is to be understood that a compound of formula (1) or a salt, solvate or *in vivo* hydrolysable ester thereof may exhibit the phenomenon of tautomerism and that the formulae drawings within this specification can represent only one of the possible tautomeric forms. It is to be understood that the invention encompasses any tautomeric form and mixtures thereof and is not to be limited merely to any one tautomeric form utilised within the formulae drawings. The formulae drawings within this specification

can represent only one of the possible tautomeric forms and it is to be understood that the specification encompasses all possible tautomeric forms of the compounds drawn not just those forms which it has been possible to show graphically herein.

It is also to be understood that certain compounds of formula (1) and salts thereof
5 can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms.

The present invention relates to the compounds of formula (1) as hereinbefore defined as well as to the salts thereof. Salts for use in pharmaceutical compositions will be pharmaceutically acceptable salts, but other salts may be useful in the production of the
10 compounds of formula (1) and their pharmaceutically acceptable salts. Pharmaceutically acceptable salts of the invention may, for example, include acid addition salts of the compounds of formula (1) as hereinbefore defined which are sufficiently basic to form such salts. Such acid addition salts include for example salts with inorganic or organic acids affording pharmaceutically acceptable anions such as with hydrogen halides (especially
15 hydrochloric or hydrobromic acid of which hydrochloric acid is particularly preferred) or with sulphuric or phosphoric acid, or with trifluoroacetic, citric or maleic acid. Suitable salts include hydrochlorides, hydrobromides, phosphates, sulphates, hydrogen sulphates, alkylsulphonates, arylsulphonates, acetates, benzoates, citrates, maleates, fumarates, succinates, lactates, tartrates, oxalates, methanesulphonates or *p*-toluenesulphonates.
20 Pharmaceutically acceptable salts of the invention may also include basic addition salts of the compounds of formula (1) as hereinbefore defined which are sufficiently acidic to form such salts. Such salts may be formed with an inorganic or organic base which affords a pharmaceutically acceptable cation. Such salts with inorganic or organic bases include for example an alkali metal salt, such as a lithium, sodium or potassium salt, an alkaline earth
25 metal salt such as a calcium or magnesium salt, an ammonium salt or an organic amine salt, for example a salt with methylamine, dimethylamine, trimethylamine, triethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine. Other basic addition salts include aluminium, zinc, benzathine, chlorprocaine, choline, diethanolamine, ethanolamine, ethyldiamine, meglumine, tromethamine or procaine.

30 The present invention further relates to an *in vivo* hydrolysable ester of a compound of formula (1). An *in vivo* hydrolysable ester of a compound of formula (1) which contains carboxy or hydroxy group is, for example a pharmaceutically acceptable ester which is cleaved in the human or animal body to produce the parent acid or alcohol. Such esters can

be identified by administering, for example, intravenously to a test animal, the compound under test and subsequently examining the test animal's body fluid.

Suitable pharmaceutically acceptable esters for carboxy include C₁₋₆alkoxymethyl esters for example methoxymethyl, C₁₋₆alkanoyloxymethyl esters for example

- 5 pivaloyloxymethyl, phthalidyl esters, C₃₋₈cycloalkoxycarbonyloxyC₁₋₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl and may be formed at any carboxy group in the compounds of this invention.

- 10 Suitable pharmaceutically-acceptable esters for hydroxy include inorganic esters such as phosphate esters (including phosphoramidic cyclic esters) and α -acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group/s. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxymethoxy. A selection of *in-vivo* hydrolysable ester forming
- 15 groups for hydroxy include C₁₋₁₀alkanoyl, for example acetyl; benzoyl; phenylacetyl; substituted benzoyl and phenylacetyl, C₁₋₁₀alkoxycarbonyl (to give alkyl carbonate esters), for example ethoxycarbonyl; di-(C₁₋₄)alkylcarbamoyl and *N*-(di-(C₁₋₄)alkylaminoethyl)-*N*-(C₁₋₄)alkylcarbamoyl (to give carbamates); di-(C₁₋₄)alkylaminoacetyl and carboxyacetyl. Examples of ring substituents on phenylacetyl and benzoyl include aminomethyl, (C₁₋
- 20 ₄)alkylaminomethyl and di-((C₁₋₄)alkyl)aminomethyl, and morpholino or piperazino linked from a ring nitrogen atom via a methylene linking group to the 3- or 4- position of the benzoyl ring. Other interesting *in-vivo* hydrolysable esters include, for example, R^AC(O)O(C₁₋₆)alkyl-CO-, wherein R^A is for example, benzyloxy-(C₁₋₄)alkyl, or phenyl). Suitable substituents on a phenyl group in such esters include, for example, 4-(C₁₋₄)piperazino-(C₁₋₄)alkyl, piperazino-
- 25 (C₁₋₄)alkyl and morpholino-(C₁₋₄)alkyl.

In this specification the term "alkyl" includes both straight-chain and branched-chain alkyl groups. However references to individual alkyl groups such as "propyl" are specific for the straight chain version only and references to individual branched-chain alkyl groups such as *t*-butyl are specific for the branched chain version only. For example,

- 30 "C₁₋₃alkyl" includes methyl, ethyl, propyl and isopropyl and examples of "C₁₋₆alkyl" include the examples of "C₁₋₃alkyl" and additionally *t*-butyl, pentyl, 2,3-dimethylpropyl, 3-methylbutyl and hexyl. Examples of "C₁₋₈alkyl" include the examples of "C₁₋₆alkyl" and additionally heptyl, 2,3-dimethylpentyl, 1-propylbutyl and octyl. An analogous convention

-7-

applies to other terms, for example "C₂₋₆alkenyl" includes vinyl, allyl, 1-propenyl, 2-butenyl, 3-butenyl, 3-methylbut-1-enyl, 1-pentenyl and 4-hexenyl and examples of "C₂₋₆alkynyl" includes ethynyl, 1-propynyl, 3-butylnyl, 2-pentylnyl and 1-methylpent-2-ynyl.

"C₃₋₇carbocyclyl" is a saturated, partially saturated or unsaturated, monocyclic ring containing 3 to 7 carbon ring atoms wherein a -CH₂- group can optionally be replaced by a -C(O)-. Suitable examples of "carbocyclyl" are cyclopropyl, cyclopentyl, cyclobutyl, cyclohexyl, cyclohexenyl, 4-oxocyclohex-1-yl and 3-oxocyclohept-5-en-1-yl.

The term "halo" refers to fluoro, chloro, bromo and iodo.

Examples of "C₁₋₆alkoxy" include methoxy, ethoxy, propoxy, isopropoxy, butyloxy, pentyloxy, 1-ethylpropoxy and hexyloxy. Examples of "C₁₋₆alkylamino" include methylamino, ethylamino, propylamino, butylamino and 2-methylpropylmino. Examples of "di(C₁₋₆alkyl)amino" include dimethylamino, *N*-methyl-*N*-ethylamino, diethylamino, *N*-propyl-*N*-3-methylbutylamino. Examples of "*N*-(C₁₋₆alkyl)-*N*-(phenyl)amino" include *N*-methyl-*N*-phenylamino, *N*-propyl-*N*-phenylamino and *N*-(2-methylbutyl)-*N*-phenylamino. Examples of "*N*-(C₁₋₆alkyl)carbamoyl" are *N*-methylcarbamoyl, *N*-ethylcarbamoyl and *N*-(2-ethylbutylcarbamoyl. Examples of "*N*-(C₁₋₆alkyl)-*N*-(phenyl)carbamoyl" include *N*-methyl-*N*-phenylcarbamoyl, *N*-butyl-*N*-phenylcarbamoyl and *N*-(3-methylpentyl)-*N*-(phenyl)carbamoyl. Examples of "*N,N*-di(C₁₋₆alkyl)carbamoyl" include *N,N*-dimethylcarbamoyl, *N*-methyl-*N*-ethylcarbamoyl and *N*-propyl-*N*-(2-methylbutyl)carbamoyl. Examples of "C₁₋₆alkylthio" include methylthio, ethylthio, propylthio, butylthio and 2-methylbutylthio.

"Heteroaryl" is a monocyclic or bicyclic aryl ring containing 5 to 10 ring atoms of which 1, 2, 3 or 4 ring atoms are chosen from nitrogen, sulphur or oxygen. Examples of heteroaryl include pyrrolyl, furanyl, thienyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, benzfuranyl, benzthieno, indolyl, benzimidazolyl, benzoxazolyl, benzthiazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, benztriazolyl, quinolinyl, isoquinolinyl and naphthiridinyl. Conveniently heteroaryl is selected from imidazolyl, pyrazolyl, thiazolyl, isoxazolyl, furanyl, thienyl, isoxazolyl, or indazolyl.

Examples of "a 3-8 membered ring optionally containing 1, 2 or 3 atoms selected from O, S and NR⁸" include oxetanyl, azetidiny, benzodiazolyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothiophenyl, tetrahydropyranyl, piperidinyl, piperazinyl, morpholinyl, homopiperidinyl and homopiperazinyl tetrahydrodioxanyl., such as oxetanyl, azetidiny, pyrrolidinyl, tetrahydrofuranyl, tetrahydropyranyl, piperidinyl, piperazinyl,

-8-

morpholinyl, homopiperidinyl and homopiperazinyl, further such as pyrrolidinyl, tetrahydropyridinyl, piperidinyl, piperazinyl, and morpholinyl.

Examples of "a 4- to 7-membered saturated heterocyclic ring system" include azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, homopiperazinyl and morpholinyl.

5 Where optional substituents are chosen from "1, 2 or 3" groups it is to be understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups. An analogous convention applies to substituents chosen from "1 or 2" groups.

10 Preferred values of R^1 , R^2 , R^3 , X and R^x are as follows. Such values may be used where appropriate with any of the definitions, claims or embodiments defined hereinbefore or hereinafter.

 In one aspect of the present invention there is provided a compound of formula (1) as depicted above wherein R^1 is C_{1-8} alkyl optionally substituted by 1, 2 or 3 substituents independently selected from nitrile, phenyl or heteroaryl, wherein phenyl and heteroaryl are 15 optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, $-OR^4$, $-SR^{10}$, C_{1-6} alkyl and trifluoromethyl.

 In another aspect of the invention R^1 is benzyl optionally substituted by 1, 2, or 3, such as 1 or 2 substituents independently selected from fluoro, chloro, bromo, methoxy, methyl and trifluoromethyl. A further aspect R^1 is benzyl.

20 In a further aspect R^1 is monofluorobenzyl such as 2-fluorobenzyl, monochlorobenzyl such as 3-chlorobenzyl, difluorobenzyl such as 2,3-difluorobenzyl, fluorochlorobenzyl such as 3-chloro-2-fluorobenzyl, trifluorobenzyl such as 2, 3, 4-trifluorobenzyl.

 In one aspect of the invention R^2 is C_{1-8} alkyl substituted by 1, 2 or 3 substituents 25 independently selected from hydroxy, amino, C_{1-6} alkoxy, C_{1-6} alkylamino, di(C_{1-6} alkyl)amino, N -(C_{1-6} alkyl)- N -(phenyl)amino, N - C_{1-6} alkylcarbamoyl, N,N -di(C_{1-6} alkyl)carbamoyl, N -(C_{1-6} alkyl)- N -(phenyl)carbamoyl, carboxy, phenoxycarbonyl, $-NR^8COR^9$, $-SO_2R^{10}$, $-SO_2NR^5R^6$ and $-NR^8SO_2R^9$.

30 In another aspect R^2 is C_{1-4} alkyl substituted by 1 or 2 hydroxy groups, conveniently 1 hydroxy group.

 In a further aspect R^2 is 2-hydroxy-1-methylethyl, 1-(hydroxymethyl)propyl, 2-hydroxy-1-(hydroxymethyl)ethyl, or 2-hydroxy-1,1-dimethylethyl, especially 2-hydroxy-1-methylethyl

-9-

In one aspect of the invention R^3 is hydrogen.

In one aspect of the invention R^4 is hydrogen, C_{1-4} alkyl for example methyl, or phenyl.

In one aspect of the invention R^5 is hydrogen, C_{1-4} alkyl or phenyl.

5 In one aspect of the invention R^6 is hydrogen, C_{1-4} alkyl or phenyl.

In one aspect of the invention R^{10} is hydrogen, C_{1-4} alkyl or phenyl.

In one aspect of the invention X is hydrogen, halo, cyano, nitro, hydroxy, thio, C_{1-6} alkylthio (optionally substituted by 1 or 2 substituents selected from halo, $-OR^{17}$, $-NR^{15}R^{16}$), C_{1-8} alkyl (optionally substituted by 1, 2 or 3 substituents independently selected from halo, -
10 OR^4 , $-NR^5R^6$, $-CONR^5R^6$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$ and $-NR^8SO_2R^9$).

In another aspect X is hydrogen, halo, cyano, nitro, hydroxy, thio, C_{1-4} alkyl or C_{1-4} alkoxy (optionally substituted by 1, 2 or 3 substituents independently selected from halo, -
15 OR^4 , $-NR^5R^6$, $-CONR^5R^6$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$ and $-NR^8SO_2R^9$), C_{1-4} alkylthio or C_{1-4} alkylamino (optionally substituted by 1 or 2 substituents selected from halo, $-OR^{17}$, $-NR^{15}R^{16}$),

In another aspect X is hydrogen, halo or C_{1-4} alkyl (optionally substituted by 1, 2 or 3 substituents independently selected from halo, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COOR^7$, -
20 NR^8COR^9 , $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$ and $-NR^8SO_2R^9$),

In another aspect X is hydrogen.

In one aspect of the invention R^x is trifluoromethyl, $-NR^5R^6$, phenyl, naphthyl, monocyclic or bicyclic heteroaryl wherein a heteroring may be partially or fully saturated and one or more ring carbon atoms may form a carbonyl group, and wherein each phenyl or heteroaryl group is optionally substituted by 1, 2 or 3 substituents independently selected
25 from halo, cyano, nitro, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COR^7$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$, $-NR^8SO_2R^9$, C_{1-6} alkyl or trifluoromethyl;
or R^x is a group selected from C_{3-7} carbocyclyl, C_{1-8} alkyl, C_{2-6} alkenyl and C_{2-6} alkynyl whereby the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COR^7$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$,
30 $-NR^8SO_2R^9$, phenyl or heteroaryl; and wherein each phenyl or heteroaryl group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COR^7$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$, $-NR^8SO_2R^9$, C_{1-6} alkyl or trifluoromethyl;

-10-

In a further aspect of the invention R^x is phenyl, heteroaryl, $-NR^5R^6$ or a group selected from C_{1-8} alkyl whereby the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$, $-NR^8SO_2R^9$, phenyl or heteroaryl; and wherein each phenyl or

5 heteroaryl group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$, $-NR^8SO_2R^9$, C_{1-6} alkyl or trifluoromethyl.

In a further aspect R^x is methyl, phenyl, 1-methylimidazolyl, 1,2-dimethylimidazolyl, or isoxazolyl.

10 In a further aspect R^x is methyl, phenyl or 1-methylimidazolyl or 1,2-dimethylimidazolyl.

In a further aspect R^x is $-NR^5R^6$ such as azetidiny, pyrrolidiny, piperaziny or morpholinyl.

A preferred class of compound is of formula (1) wherein;

15 R^1 is C_{1-8} alkyl optionally substituted by 1, 2 or 3 substituents independently selected from nitrile, phenyl or heteroaryl, wherein phenyl and heteroaryl are optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, $-OR^4$, $-SR^{10}$, C_{1-6} alkyl and trifluoromethyl;

R^2 is C_{1-8} alkyl substituted by 1, 2 or 3 substituents independently selected from hydroxy, amino, C_{1-6} alkoxy, C_{1-6} alkylamino, di(C_{1-6} alkyl)amino, N -(C_{1-6} alkyl)- N -(phenyl)amino, N - C_{1-6} alkylcarbamoyl, N,N -di(C_{1-6} alkyl)carbamoyl, N -(C_{1-6} alkyl)- N -(phenyl)carbamoyl, carboxy, phenoxycarbonyl, $-NR^8COR^9$, $-SO_2R^{10}$, $-SO_2NR^5R^6$ and $-NR^8SO_2R^9$;

20

R^3 is hydrogen;

$R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}$ and R^{17} are independently hydrogen, C_{1-4} alkyl or phenyl; and

25

X is hydrogen, halo, cyano, nitro, hydroxy, thio, C_{1-6} alkylthio (optionally substituted by 1 or 2 substituents selected from halo, $-OR^{17}$, $-NR^{15}R^{16}$), C_{1-8} alkyl (optionally substituted by 1, 2 or 3 substituents independently selected from halo, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$ and $-NR^8SO_2R^9$);

30 R^x is $-NR^5R^6$, phenyl, naphthyl, monocyclic or bicyclic heteroaryl wherein a heteroring may be partially or fully saturated and one or more ring carbon atoms may form a carbonyl group, and wherein each phenyl or heteroaryl group is optionally substituted by 1, 2 or 3 substituents

-11-

independently selected from halo, cyano, nitro, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COR^7$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$, $-NR^8SO_2R^9$, C_{1-6} alkyl or trifluoromethyl;

or R^x is C_{1-8} alkyl optionally substituted by 1, 2 or 3 substituents independently selected from halo, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COR^7$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$,

- 5 $-NR^8SO_2R^9$, phenyl or heteroaryl; and wherein each phenyl or heteroaryl group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COR^7$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$, $-NR^8SO_2R^9$, C_{1-6} alkyl or trifluoromethyl;

Another preferred class of compound is of formula (1) wherein;

- 10 R^1 is benzyl optionally substituted by 1, 2 or 3, such as 2, substituents independently selected from fluoro, chloro, bromo, methoxy, methyl and trifluoromethyl;

R^2 is C_{1-4} alkyl substituted by 1 or 2 hydroxy groups;

R^3 is hydrogen;

X is hydrogen; and

- 15 R^x is methyl, phenyl, 1-methylimidazolyl, 1,2-dimethylimidazolyl, isoxazolyl or *N,N*-dimethylamino. Alternatively R^x is $-NR^5R^6$ -such as azetidiny, pyrrolidinyl piperazinyl, piperidinyl or morpholinyl

In another class R^1 is benzyl optionally substituted by 3-chloro-2-fluoro, 2,3-difluoro or 2,3,4-trifluoro

- 20 R^2 is 2-hydroxy-1-methylethyl

R^3 is hydrogen

X is hydrogen

R^x is azetidiny, pyrrolidinyl or morpholinyl, *N,N*-dimethylamino, piperidinyl, methyl, 1-methylimidazolyl and 1,2-dimethylimidazolyl.

- 25 Convenient compounds of the invention include each exemplified compound, each selected independently and pharmaceutically acceptable salts, in vivo hydrolysable esters thereof.

Particular compounds of the invention include:

N-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino})-pyrimidin-4-yl)methanesulfonamide

- 30

N-[2-[(3-Chloro-2-fluorobenzyl)thio]-6-[(2-hydroxy-1-methylethyl)amino]-4-pyrimidinyl]-4-morpholinesulfonamide

-12-

N-[2-[[[3-Chloro-2-fluorophenyl)methyl]thio]-6-[(2-hydroxy-1-methylethyl)amino]-4-pyrimidinyl]-1,2-dimethyl-1H-imidazole-4-sulfonamide

N-(2-[(2,3-Difluorobenzyl)thio]-6-[[[(1R)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)piperidine-1-sulfonamide

5 N-(2-[(2,3-Difluorobenzyl)thio]-6-[[[(1R)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)pyrrolidine-1-sulfonamide

N-(2-[(2,3-Difluorobenzyl)thio]-6-[[[(1R)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)azetidine-1-sulfonamide

10 N-{6-[[[(1R)-2-Hydroxy-1-methylethyl]amino}-2-[(2,3,4-trifluorobenzyl)thio]-pyrimidin-4-yl}morpholine-4-sulfonamide

N-(2-[(2,3-Difluorobenzyl)thio]-6-[[[(1R)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)morpholine-4-sulfonamide

N-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[[[(1R)-2-hydroxy-1-methylethyl]amino}-pyrimidin-4-yl)azetidine-1-sulfonamide

15 N-{6-[[[(1R)-2-Hydroxy-1-methylethyl]amino}-2-[(2,3,4-trifluorobenzyl)thio]-pyrimidin-4-yl}azetidine-1-sulfonamide

N-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[[[(1R)-2-hydroxy-1-methylethyl]amino}-pyrimidin-4-yl)-N,N-dimethylsulfamide, and

20 N-[2-[[[3-Chloro-2-fluorophenyl)methyl]thio]-6-[(R)-(2-hydroxy-1-methylethyl)amino]-4-pyrimidinyl]-1-methyl-1H-imidazole-4-sulfonamide

and pharmaceutically acceptable salts, solvates or *in vivo* hydrolysable esters thereof. Each of the above mentioned compound and the pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, individually is a preferred aspect of the invention.

Further particular compounds of the invention include:

25 N-(2-[(2,3-difluorobenzyl)thio]-6-[[[(1R)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)methanesulfonamide;

N-(2-[(2,3-difluorobenzyl)thio]-6-[[[(1R)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-1-methyl-1H-imidazole-4-sulfonamide;

N-(2-(benzylthio)-6-[[[(1R)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-

30 methanesulfonamide; and

N-(2-(benzylthio)-6-[[[(1R)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)benzenesulfonamide;


-13-

and pharmaceutically acceptable salts, solvates or *in vivo* hydrolysable esters thereof. Each of the above mentioned compound and the pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, individually is a preferred aspect of the invention.

The present invention further provides processes for the preparation of compounds
5 of formula (1) as defined above which comprise:

Process 1

(a) treating a compound of formula (2):



10 (2)

wherein R¹, R² and R³ are as defined in formula (1) and X is hydrogen with sulfonyl chlorides (R^xSO₂Cl) where R^x is as defined in formula (1).


and optionally thereafter (i), (ii), (iii), (iv), or (v) in any order:

- 15 i) removing any protecting groups;
ii) converting the compound of formula (1) into a further compound of formula (1)
iii) forming a salt
iv) forming a prodrug
v) forming an *in vivo* hydrolysable ester.

20 Reaction of compounds of formula (2) wherein R¹, R² and R³ are as defined in formula (1) and X is hydrogen with sulfonyl chlorides (R^xSO₂Cl) can be carried out in the presence of a suitable base and solvent. Examples of suitable bases include trialkylamines, such as triethylamine or *N,N*-diisopropylethylamine or pyridine (optionally in the presence of a catalyst such as 4-dimethylaminopyridine). Suitable solvents include dichloromethane,
25 pyridine, *N,N*-dimethylamides, 1-methyl-2-pyrrolidone, and ethers such as tetrahydrofuran, 1,4-dioxane, glyme and diglyme. Preferably *N,N*-dimethylformamide is used. The temperature of the reaction can be performed between -10°C and 100°C. Preferably *N,N*-diisopropylethylamine in dichloromethane or pyridine with 4-dimethylaminopyridine both at ambient temperature are used.

-14-

Compounds of formula (2) wherein R^1 , R^2 and R^3 and X are as defined in formula (1), can be prepared from compounds of formula (3) wherein R^1 and X are as defined in formula (1) and L is halogen by treatment with nucleophilic amines NR^2R^3 as defined in formula (1) in the presence of a suitable base and solvent.




(3)

Examples of suitable bases include trialkylamines, such as triethylamine or *N,N*-diisopropylethylamine. Suitable solvents include *N,N*-dimethylamides, 1-methyl-2-pyrrolidone, and ethers such as tetrahydrofuran, 1,4-dioxane, glyme and diglyme. The temperature of the reaction can be performed between 0°C and 150°C. Preferably *N,N*-diisopropylethylamine in 1-methyl-2-pyrrolidinone at 120°C is used.

Compounds of formula (3) wherein R^1 and X are as defined in formula (1) and L is halogen may be prepared by treating a compound of formula (3) wherein R^1 and X are as defined in formula (1) and L is OH with a halogenating agent such as phosphorous oxychloride. The reaction may be carried out in the presence of an *N,N*-dialkylaniline, such as *N,N*-dimethylaniline at reflux.

Compounds of formula (3) wherein R^1 and X are as defined in formula (1) and L is OH;



(4)


may be prepared from compounds of formula (4) wherein X is as defined in formula (1) by reaction with alkylhalides (R_1A) where R_1 is as defined in formula (1) and A is halogen in the presence of a suitable base and solvent.

Examples of suitable bases include the alkali metal hydroxides such as Li, Na, or K, or metal carbonates such as Li, Na, K or Cs, or metal acetates such as Li, Na, K or Cs, or metal alkoxides such as Li, Na, K-*tert*-butoxide. Suitable solvents include *N,N*-dimethylamides, 1-methyl-2-pyrrolidinone, ethers such as tetrahydrofuran, 1,4-dioxane, glyme

-15-

and diglyme and alcohols such as methanol, ethanol and *tert*-butanol. Preferably potassium hydroxide in *N,N*-dimethylformamide at ambient temperature is used.


Compounds of formula (3) wherein R^1 and X are as defined in formula (1) and L is halogen;



may also be prepared from compounds of formula (5) wherein X and R^1 are as defined in formula (1) and L is halogen by reaction with concentrated ammonium hydroxide solution in the presence of a suitable solvent. Suitable solvents include *N*-methyl-2-pyrrolidone, acetonitrile and ethers such as tetrahydrofuran, 1,4-dioxane, glyme and diglyme. The temperature of the reaction can be performed between 0°C and 150°C. Preferably acetonitrile at 60°C is used.

Compounds of formula (5) wherein R^1 and X are as defined in formula (1) and L is halogen may be prepared from compounds of formula (5) wherein R^1 and X are as defined in formula (1) and L is OH by reaction with a halogenating agent such as phosphorous oxychloride. The reaction may be carried out in the presence of *N,N*-dimethylaniline at reflux.

Compounds of formula (5) wherein R^1 and X are as defined in formula (1) and L is OH;



may be prepared from compounds of formula (6) wherein X are as defined in formula (1) and L is OH by reaction with alkylhalides (R_1A) where R_1 is as defined in formula (1) and A is halogen in the presence of a suitable base and solvent.

25 Examples of suitable bases include the alkali metal hydroxides such as Li, Na, or K, or metal carbonates such as Li, Na, K or Cs, or metal acetates such as Li, Na, K or Cs, or metal alkoxides such as Li, Na, K *tert*-butoxide. Suitable solvents include *N,N*-dimethylamides, 1-methyl-2-pyrrolidone, ethers such as tetrahydrofuran, 1,4-dioxane, glyme

-16-


and diglyme and alcohols such as methanol, ethanol and *tert*-butanol. Preferably potassium hydroxide in *N,N*-dimethylformamide at ambient temperature is used.

Compounds of formulae (4) and (6) are either commercially available, are well known in the literature or may be easily prepared using known techniques.

5

Process 2

(b) treating a compound of formula (7):



(7)

10

wherein R¹, R^x and X are as defined in formula (1), L is a halogen and Y is either hydrogen or a protecting group with nucleophilic amines of the type NR²R³ as defined in formula (1) in the presence or absence of a suitable base and solvent.

and optionally thereafter (i), (ii), (iii), (iv) or (v) in any order:

- 15 i) removing any protecting groups;
ii) converting the compound of formula (1) into a further compound of formula (1)
iii) forming a salt
iv) forming a prodrug
v) forming an *in vivo* hydrolysable ester.


20

Examples of suitable bases include trialkylamines, such as triethylamine or *N,N*-diisopropylethylamine. Suitable solvents include *N,N*-dimethylamides, 1-methyl-2-pyrrolidone, and ethers such as tetrahydrofuran, 1,4-dioxane, glyme and diglyme. The temperature of the reaction can be performed between 0°C and 150°C. Preferably 1-methyl-2-pyrrolidinone at 80°C is used.

25

Compounds of formula (7) wherein R¹, R^x and X are as defined in formula (1) and L is halogen and Y is a protecting group or hydrogen;

-17-



(5)

may be prepared from compounds of formula (5) wherein X and R¹ are as defined in formula (1) and L is a halogen by reaction with sulfonamides or sulfamides of formula R^xSO₂NH₂

5 where R^x is as defined in formula (1) in the presence of a suitable base and solvent.

Examples of suitable bases include the alkali metal hydrides such as Na or K.

Suitable solvents include *N,N*-dimethylamides, 1-methyl-2-pyrrolidinone, ethers such as tetrahydrofuran, 1,4-dioxane, glyme and diglyme. The temperature of the reaction may be performed between 0°C and 100°C. Preferably sodium hydride in *N,N*-dimethylformamide at

10 ambient temperature is employed.

Compounds of formula (5) can be prepared as described in process (1).


Compounds of formula R^xSO₂NH₂ where R^x is NR⁵R⁶ may be prepared from sulfamide in the presence of a suitable solvent. Suitable solvents include *N,N*-

dimethylamides, 1-methyl-2-pyrrolidinone, dichloromethane, chloroform, ethers such as

15 tetrahydrofuran, 1,4-dioxane, glyme and diglyme and alcohols such as methanol, ethanol and *tert*-butanol. Preferably 1,4-dioxane is used at 110°C.

Process 3

(c) treating a compound of formula (8):



20

(8)

wherein R¹, R², R³, R^x and X are as defined in formula (1) and L is halogen, with

sulfonamides of formula R^xSO₂NH₂ where R^x is as defined in formula (1) except NR⁵R⁶ in

25 the presence of a suitable base and solvent.

and optionally thereafter (i), (ii), (iii), (iv) or (v) in any order:

i) removing any protecting groups;

-18-


- ii) converting the compound of formula (1) into a further compound of formula (1)
- iii) forming a salt
- iv) forming a prodrug
- v) forming an *in vivo* hydrolysable ester.

5

Examples of suitable bases include the alkali metal hydrides such as Na or K, or metal alkoxides such as Li, Na or K-*tert*-butoxide, or metal carbonates such as Na, K, Cs. Suitable solvents include *N,N*-dimethylamides, 1-methyl-2-pyrrolidinone, ethers such as tetrahydrofuran, 1,4-dioxane, glyme and diglyme. Preferably sodium hydride in *N,N*-

10 dimethylformamide at ambient temperature is employed.

Compounds of formula (8) wherein R^1 , R^2 , R^3 and X are as defined in formula (1) and L is halogen;



(5)

- 15 may be prepared from compounds of formula (5) wherein X and R^1 are as defined in formula (1) and L is a halogen by reaction with nucleophilic amines NR^2R^3 as defined in formula (1) in the presence a suitable base and solvent. Examples of suitable bases include trialkylamines, such as triethylamine or *N,N*-diisopropylethylamine. Suitable solvents include *N,N*-dimethylamides, 1-methyl-2-pyrrolidone, and ethers such as tetrahydrofuran, 1,4-dioxane, glyme and diglyme. The temperature of the reaction can be performed between 0°C and 150°C. Preferably *N,N*-diisopropylethylamine in *N*-methylpyrrolidinone is used at room temperature.

Compounds of formula (5) can be prepared as described in process (1).

- Compounds of formula $R^xSO_2NH_2$ where R^x is as defined in formula (1), except
25 NR^5R^6 , are either commercially available or well known in the literature or may be prepared from the corresponding commercially available or well known in the literature sulfonyl chlorides R^xSO_2Cl .

It will be appreciated by those skilled in the art that in the processes of the present invention certain functional groups such as hydroxyl or amino groups in the starting reagents or
30 intermediate compounds may need to be protected by protecting groups. Thus, the

-19-

preparation of the compounds of formula (1) may involve, at an appropriate stage, the removal of one or more protecting groups. The protection and deprotection of functional groups is fully described in 'Protective Groups in Organic Chemistry', edited by J. W. F. McOmie, Plenum Press (1973), and 'Protective Groups in Organic Synthesis', 2nd edition, T.

5 W. Greene & P. G. M. Wuts, Wiley-Interscience (1991).

Compounds of formulae (2), (3), (4) and (5), (6), (7), and (8) are either commercially available, are well known in the literature or may be easily prepared using known techniques.

A compound of formula (1) may be prepared from another compound of formula
10 (1) by chemical modification. Examples of chemical modifications include standard alkylation, arylation, heteroarylation, acylation, sulphonylation, phosphorylation, aromatic halogenation and coupling reactions. These reactions may be used to add new substituents or to modify existing substituents. Alternatively, existing substituents in compounds of formula (1) may be modified by, for example, oxidation, reduction, elimination, hydrolysis or other
15 cleavage reactions to yield other compounds of formula (1).

Novel intermediate compounds form a further aspect of the invention.

The compounds of formula (1) above may be converted to a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, as discussed above. The salt is preferably a basic addition salt.

20 The compounds of formula (1) have activity as pharmaceuticals, in particular as modulators of chemokine receptor (especially CXCR2) activity, and may be used in the treatment (therapeutic or prophylactic) of conditions/diseases in human and non-human animals which are exacerbated or caused by excessive or unregulated production of chemokines. Examples of such conditions/diseases include (each taken independently):

25

- (1) **(the respiratory tract)** obstructive airways diseases including chronic obstructive pulmonary disease (COPD); asthma, such as bronchial, allergic, intrinsic, extrinsic and dust asthma, particularly chronic or inveterate asthma (e.g. late asthma and airways hyper-responsiveness); bronchitis; acute, allergic,
30 atrophic rhinitis and chronic rhinitis including rhinitis caseosa, hypertrophic rhinitis, rhinitis purulenta, rhinitis sicca and rhinitis medicamentosa; membranous rhinitis including croupous, fibrinous and pseudomembranous rhinitis and scrofulous rhinitis; seasonal rhinitis including rhinitis nervosa (hay fever) and

-20-

vasomotor rhinitis; sarcoidosis, farmer's lung and related diseases, fibroid lung and idiopathic interstitial pneumonia;

- 5 (2) **(bone and joints)** rheumatoid arthritis, seronegative spondyloarthropathies (including ankylosing spondylitis, psoriatic arthritis and Reiter's disease), Behchet's disease, Sjogren's syndrome and systemic sclerosis;
- 10 (3) **(skin)** psoriasis, atopic dermatitis, contact dermatitis and other eczmatous dermitides, seborrhoetic dermatitis, Lichen planus, Pemphigus, bullous Pemphigus, Epidermolysis bullosa, urticaria, angiodermas, vasculitides, erythemas, cutaneous eosinophilias, uveitis, Alopecia areata and vernal conjunctivitis;
- 15 (4) **(gastrointestinal tract)** Coeliac disease, proctitis, eosinophilic gastro-enteritis, mastocytosis, Crohn's disease, ulcerative colitis, indeterminate colitis, microscopic colitis, inflammatory bowel disease, irritable bowel syndrome, non-inflammatory diarrhea, food-related allergies which have effects remote from the gut, e.g., migraine, rhinitis and eczema;
- 20 (5) **(central and peripheral nervous system)** Neurodegenerative diseases and dementia disorders, e.g. Alzheimer's disease, amyotrophic lateral sclerosis and other motor neuron diseases, Creutzfeldt-Jacob's disease and other prion diseases, HIV encephalopathy (AIDS dementia complex), Huntington's disease, frontotemporal dementia, Lewy body dementia and vascular dementia;
- 25 polyneuropathies, e.g. Guillain-Barré syndrome, chronic inflammatory demyelinating polyradiculoneuropathy, multifocal motor neuropathy, plexopathies; CNS demyelination, e.g. multiple sclerosis, acute disseminated/haemorrhagic encephalomyelitis, and subacute sclerosing panencephalitis; neuromuscular disorders, e.g. myasthenia gravis and Lambert-
- 30 Eaton syndrome; spinal disorders, e.g. tropical spastic paraparesis, and stiff-man syndrome; paraneoplastic syndromes, e.g. cerebellar degeneration and encephalomyelitis; CNS trauma; migraine; and stroke.

-21-

- (6) **(other tissues and systemic disease)** atherosclerosis, Acquired Immunodeficiency Syndrome (AIDS), lupus erythematosus, systemic lupus, erythematosus, Hashimoto's thyroiditis, type I diabetes, nephrotic syndrome, eosinophilia fascitis, hyper IgE syndrome, lepromatous leprosy, and idiopathic thrombocytopenia pupura; post-operative adhesions, and sepsis.
- (7) **(allograft rejection)** acute and chronic following, for example, transplantation of kidney, heart, liver, lung, bone marrow, skin and cornea; and chronic graft versus host disease;
- (8) Cancers, especially non-small cell lung cancer (NSCLC), malignant melanoma, prostate cancer and squamous sarcoma, and tumour metastasis, non melanoma skin cancer and chemoprevention metastases;
- (9) Diseases in which angiogenesis is associated with raised CXCR2 chemokine levels (e.g. NSCLC, diabetic retinopathy);
- (10) Cystic fibrosis;
- (11) Burn wounds & chronic skin ulcers;
- (12) Reproductive Diseases (e.g. Disorders of ovulation, menstruation and implantation, Pre-term labour, Endometriosis);
- (13) Re-perfusion injury in the heart, brain, peripheral limbs and other organs, inhibition of atherosclerosis.

Thus, the present invention provides a compound of formula (1), or a pharmaceutically-acceptable salt, solvate or an *in vivo* hydrolysable ester thereof, as hereinbefore defined for use in therapy.

Preferably the compounds of the invention are used to treat diseases in which the chemokine receptor belongs to the CXC chemokine receptor subfamily, more preferably the target chemokine receptor is the CXCR2 receptor.

Particular conditions which can be treated with the compounds of the invention are cancer, diseases in which angiogenesis is associated with raised CXCR2 chemokine levels, and inflammatory diseases such as asthma, allergic rhinitis, COPD, rheumatoid arthritis, psoriasis, inflammatory bowel diseases, osteoarthritis or osteoporosis.

5 As a further aspect of the present invention, certain compounds of formula (1) may have utility as antagonists of the CX3CR1 receptor. Such compounds are expected to be particularly useful in the treatment of disorders within the central and peripheral nervous system and other conditions characterized by an activation of microglia and/or infiltration of leukocytes (e.g. stroke/ischemia and head trauma).

10 In a further aspect, the present invention provides a compound of formula (1), or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, as hereinbefore defined for use as a medicament.

 In a still further aspect, the present invention provides the use of a compound of formula (1), or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester
15 thereof, as hereinbefore defined for use as a medicament for the treatment of human diseases or conditions in which modulation of chemokine receptor activity is beneficial.

 In a still further aspect, the present invention provides the use of a compound of formula (1), or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, as hereinbefore defined for use as a medicament for the treatment of asthma, allergic
20 rhinitis, cancer, COPD, rheumatoid arthritis, psoriasis, inflammatory bowel diseases, osteoarthritis or osteoporosis.

 In a further aspect, the present invention provides the use of a compound of formula (1), or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, as hereinbefore defined in the manufacture of a medicament for use in therapy.

25 In a still further aspect, the present invention provides the use of a compound of formula (1), or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, as hereinbefore defined in the manufacture of a medicament for the treatment of human diseases or conditions in which modulation of chemokine receptor activity is beneficial.

30 In a still further aspect, the present invention provides the use of a compound of formula (1), or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, as hereinbefore defined in the manufacture of a medicament for the treatment of

-23-

asthma, allergic rhinitis, cancer, COPD, rheumatoid arthritis, psoriasis, inflammatory bowel diseases, osteoarthritis or osteoporosis.

In the context of the present specification, the term "therapy" also includes "prophylaxis" unless there are specific indications to the contrary. The terms "therapeutic" and "therapeutically" should be construed accordingly.

The invention still further provides a method of treating a chemokine mediated disease wherein the chemokine binds to a chemokine (especially CXCR2) receptor, which comprises administering to a patient a therapeutically effective amount of a compound of formula , or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester, as hereinbefore defined.

The invention also provides a method of treating an inflammatory disease, especially asthma, allergic rhinitis, COPD, rheumatoid arthritis, psoriasis, inflammatory bowel diseases, osteoarthritis or osteoporosis, in a patient suffering from, or at risk of, said disease, which comprises administering to the patient a therapeutically effective amount of a compound of formula (1), or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, as hereinbefore defined.

For the above-mentioned therapeutic uses the dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the disorder indicated.

The compounds of formula (1) and pharmaceutically acceptable salts, solvates or *in vivo* hydrolysable esters thereof may be used on their own but will generally be administered in the form of a pharmaceutical composition in which formula (1) compound/salt/solvate/ester (active ingredient) is in association with a pharmaceutically acceptable adjuvant, diluent or carrier. Depending on the mode of administration, the pharmaceutical composition will preferably comprise from 0.05 to 99 %w (per cent by weight), more preferably from 0.05 to 80 %w, still more preferably from 0.10 to 70 %w, and even more preferably from 0.10 to 50 %w, of active ingredient, all percentages by weight being based on total composition.

The present invention also provides a pharmaceutical composition comprising a compound of formula (1), or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, as hereinbefore defined, in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

-24-

The invention further provides a process for the preparation of a pharmaceutical composition of the invention which comprises mixing a compound of formula (1), or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, as hereinbefore defined, with a pharmaceutically acceptable adjuvant, diluent or carrier. The pharmaceutical compositions may be administered topically (e.g. to the lung and/or airways or to the skin) in the form of solutions, suspensions, heptafluoroalkane aerosols and dry powder formulations; or systemically, e.g. by oral administration in the form of tablets, capsules, syrups, powders or granules, or by parenteral administration in the form of solutions or suspensions, or by subcutaneous administration or by rectal administration in the form of suppositories or transdermally. Preferably the compounds of the invention are administered orally.

In addition to their use as therapeutic medicines, the compounds of formula (1) and their pharmaceutically acceptable salts, solvate or *in vivo* hydrolysable esters are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effect of chemokine modulation activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

The invention further relates to combination therapies wherein a compound of formula (1) or a pharmaceutically acceptable salts, solvate or *in vivo* hydrolysable ester thereof, or a pharmaceutical composition or formulation comprising a compound of formula (1) is administered concurrently or sequentially with therapy and/or an agent for the treatment of any one of asthma, allergic rhinitis, cancer, COPD, rheumatoid arthritis, psoriasis, inflammatory bowel disease, irritable bowel syndrome, osteoarthritis or osteoporosis.

In particular, for the treatment of the inflammatory diseases rheumatoid arthritis, psoriasis, inflammatory bowel disease, irritable bowel syndrome, COPD, asthma and allergic rhinitis the compounds of the invention may be combined with agents such as TNF- α inhibitors such as anti-TNF monoclonal antibodies (such as Remicade, CDP-870 and D.sub2.E.sub7.) and TNF receptor immunoglobulin molecules (such as Enbrel.reg.), non-selective COX-1 / COX-2 inhibitors (such as piroxicam, diclofenac, propionic acids such as naproxen, flubiprofen, fenoprofen, ketoprofen and ibuprofen, fenamates such as mefenamic acid, indomethacin, sulindac, apazone, pyrazolones such as phenylbutazone, salicylates such as aspirin), COX-2 inhibitors (such as meloxicam, celecoxib, rofecoxib, valdecoxib and etoricoxib) low dose methotrexate, lefunomide; ciclesonide; hydroxychloroquine, d-penicillamine, auranofin or parenteral or oral gold. For inflammatory bowel disease and

-25-

irritable bowel disorder further convenient agents include sulphasalazine and 5-ASAs, topical and systemic steroids, immunomodulators and immunosuppressants, antibiotics, probiotics and anti-integrins.

The present invention still further relates to the combination of a compound of the
5 invention together with a leukotriene biosynthesis inhibitor, 5-lipoxygenase (5-LO) inhibitor or 5-lipoxygenase activating protein (FLAP) antagonist such as zileuton; ABT-761; fenleuton; tepoxalin; Abbott-79175; Abbott-85761; N-(5-substituted)-thiophene-2-alkylsulfonamides; 2,6-di-tert-butylphenol hydrazones; methoxytetrahydropyrans such as Zeneca ZD-2138; the compound SB-210661; pyridinyl-substituted 2-cyanonaphthalene compounds such as L-
10 739,010; 2-cyanoquinoline compounds such as L-746,530; indole and quinoline compounds such as MK-591, MK-886, and BAY x 1005.

The present invention still further relates to the combination of a compound of the invention together with a receptor antagonist for leukotrienes LTB.sub4., LTC.sub4., LTD.sub4., and LTE.sub4. selected from the group consisting of the phenothiazin-3-ones
15 such as L-651,392; amidino compounds such as CGS-25019c; benzoxalamines such as ontazolast; benzenecarboximidamides such as BIIL 284/260; and compounds such as zafirlukast, ablukast, montelukast, pranlukast, verlukast (MK-679), RG-12525, Ro-245913, iralukast (CGP 45715A), and BAY x 7195.

The present invention still further relates to the combination of a compound of the
20 invention together with a PDE4 inhibitor including inhibitors of the isoform PDE4D.

The present invention still further relates to the combination of a compound of the invention together with a antihistaminic H.sub1. receptor antagonists such as cetirizine, loratadine, desloratadine, fexofenadine, astemizole, azelastine, and chlorpheniramine.

The present invention still further relates to the combination of a compound of the
25 invention together with a gastroprotective H.sub2. receptor antagonist.

The present invention still further relates to the combination of a compound of the invention together with an α .sub1.- and α .sub2.-adrenoceptor agonist vasoconstrictor sympathomimetic agent, such as propylhexedrine, phenylephrine, phenylpropanolamine, pseudoephedrine, naphazoline hydrochloride, oxymetazoline hydrochloride, tetrahydrozoline
30 hydrochloride, xylometazoline hydrochloride, and ethylnorepinephrine hydrochloride.

-26-

The present invention still further relates to the combination of a compound of the invention together with anticholinergic agents such as ipratropium bromide; tiotropium bromide; oxitropium bromide; pirenzepine; and telenzepine.

The present invention still further relates to the combination of a compound of the invention together with a β .sub1.- to β .sub4.-adrenoceptor agonists such as metaproterenol, isoproterenol, isoprenaline, albuterol, salbutamol, formoterol, salmeterol, terbutaline, orciprenaline, bitolterol mesylate, and pirbuterol; or methylxanthanines including theophylline and aminophylline; sodium cromoglycate; or muscarinic receptor (M1, M2, and M3) antagonist.

10 The present invention still further relates to the combination of a compound of the invention together with an insulin-like growth factor type I (IGF-1) mimetic.

The present invention still further relates to the combination of a compound of the invention together with an inhaled glucocorticoid with reduced systemic side effects, such as prednisone, prednisolone, flunisolide, triamcinolone acetonide, beclomethasone dipropionate, 15 budesonide, fluticasone propionate, and mometasone furoate.

The present invention still further relates to the combination of a compound of the invention together with an inhibitor of matrix metalloproteases (MMPs), i.e., the stromelysins, the collagenases, and the gelatinases, as well as aggrecanase; especially collagenase-1 (MMP-1), collagenase-2 (MMP-8), collagenase-3 (MMP-13), stromelysin-1 (MMP-3), stromelysin-2 20 (MMP-10), and stromelysin-3 (MMP-11) and MMP-12.

The present invention still further relates to the combination of a compound of the invention together with other modulators of chemokine receptor function such as CCR1, CCR2, CCR2A, CCR2B, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CCR10 and CCR11 (for the C-C family); CXCR1, CXCR3, CXCR4 and CXCR5 (for the C-X-C family) 25 and CX₃CR1 for the C-X₃-C family.

The present invention still further relates to the combination of a compound of the invention together with antiviral agents such as Viracept, AZT, aciclovir and famciclovir, and antiseptis compounds such as Valant.

The present invention still further relates to the combination of a compound of the invention together with cardiovascular agents such as calcium channel blockers, lipid lowering agents such as statins, fibrates, beta-blockers, Ace inhibitors, Angiotensin-2 receptor antagonists and platelet aggregation inhibitors.

-27-

The present invention still further relates to the combination of a compound of the invention together with CNS agents such as antidepressants (such as sertraline), anti-Parkinsonian drugs (such as deprenyl, L-dopa, Requip, Mirapex, MAOB inhibitors such as selegine and rasagiline, COMT inhibitors such as Tasmar, A-2 inhibitors, dopamine reuptake
5 inhibitors, NMDA antagonists, Nicotine agonists, Dopamine agonists and inhibitors of neuronal nitric oxide synthase), and anti-Alzheimer's drugs such as donepezil, tacrine, COX-2 inhibitors, propentofylline or metryfonate.

The present invention still further relates to the combination of a compound of the invention together with (i) tryptase inhibitors; (ii) platelet activating factor (PAF) antagonists;
10 (iii) interleukin converting enzyme (ICE) inhibitors; (iv) IMPDH inhibitors; (v) adhesion molecule inhibitors including VLA-4 antagonists; (vi) cathepsins; (vii) MAP kinase inhibitors; (viii) glucose-6 phosphate dehydrogenase inhibitors; (ix) kinin-B.sub1. - and B.sub2. -receptor antagonists; (x) anti-gout agents, e.g., colchicine; (xi) xanthine oxidase inhibitors, e.g., allopurinol; (xii) uricosuric agents, e.g., probenecid, sulfinpyrazone, and
15 benzbromarone; (xiii) growth hormone secretagogues; (xiv) transforming growth factor (TGF β); (xv) platelet-derived growth factor (PDGF); (xvi) fibroblast growth factor, e.g., basic fibroblast growth factor (bFGF); (xvii) granulocyte macrophage colony stimulating factor (GM-CSF); (xviii) capsaicin cream; (xix) Tachykinin NK.sub1. and NK.sub3. receptor antagonists selected from the group consisting of NKP-608C; SB-233412 (talnetant); and D-
20 4418; (xx) elastase inhibitors selected from the group consisting of UT-77 and ZD-0892; (xxi) TNF δ converting enzyme inhibitors (TACE); (xxii) induced nitric oxide synthase inhibitors (iNOS) or (xxiii) chemoattractant receptor-homologous molecule expressed on TH2 cells, (CRTH2 antagonists).

The compounds of the present invention may also be used in combination with
25 osteoporosis agents such as roloxifene, droloxifene, lasofoxifene or fosomax and immunosuppressant agents such as FK-506, rapamycin, cyclosporine, azathioprine, and methotrexate;.

The compounds of the invention may also be used in combination with existing therapeutic agents for the treatment of osteoarthritis. Suitable agents to be used in
30 combination include standard non-steroidal anti-inflammatory agents (hereinafter NSAID's) such as piroxicam, diclofenac, propionic acids such as naproxen, flubiprofen, fenoprofen, ketoprofen and ibuprofen, fenamates such as mefenamic acid, indomethacin, sulindac, apazone, pyrazolones such as phenylbutazone, salicylates such as aspirin, COX-2 inhibitors

-28-

such as celecoxib, valdecoxib, rofecoxib and etoricoxib, analgesics and intraarticular therapies such as corticosteroids and hyaluronic acids such as hyalgan and synvisc and P2X7 receptor antagonists.

The compounds of the invention can also be used in combination with existing

5 therapeutic agents for the treatment of cancer. Suitable agents to be used in combination include:

- (i) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as alkylating agents (for example cis-platin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulphan and nitrosoureas); antimetabolites (for
10 example antifolates such as fluoropyrimidines like 5-fluorouracil and tegafur, raltitrexed, methotrexate, cytosine arabinoside, hydroxyurea, gemcitabine and paclitaxel (Taxol®); antitumour antibiotics (for example anthracyclines like adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin and mithramycin); antimitotic agents (for example vinca alkaloids like vincristine, vinblastine, vindesine and
15 vinorelbine and taxoids like taxol and taxotere); and topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan and camptothecin);
- (ii) cytostatic agents such as antioestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene and idoxifyfene), oestrogen receptor down regulators (for example fulvestrant), antiandrogens (for example bicalutamide, flutamide, nilutamide and cyproterone acetate),
20 LHRH antagonists or LHRH agonists (for example goserelin, leuporelin and buserelin), progestogens (for example megestrol acetate), aromatase inhibitors (for example as anastrozole, letrozole, vorazole and exemestane) and inhibitors of 5 α -reductase such as finasteride;
- (iii) Agents which inhibit cancer cell invasion (for example metalloproteinase inhibitors like
25 marimastat and inhibitors of urokinase plasminogen activator receptor function);
- (iv) inhibitors of growth factor function, for example such inhibitors include growth factor antibodies, growth factor receptor antibodies (for example the anti-erb2 antibody trastuzumab [Herceptin™] and the anti-erb1 antibody cetuximab [C225]), farnesyl transferase inhibitors, tyrosine kinase inhibitors and serine/threonine kinase inhibitors, for
30 example inhibitors of the epidermal growth factor family (for example EGFR family tyrosine kinase inhibitors such as N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, AZD1839), N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) and 6-acrylamido-N-(3-chloro-

-29-

4-fluorophenyl)-7-(3-morpholinopropoxy)quinazolin-4-amine (CI 1033)), for example inhibitors of the platelet-derived growth factor family and for example inhibitors of the hepatocyte growth factor family;

- (v) antiangiogenic agents such as those which inhibit the effects of vascular endothelial growth factor, (for example the anti-vascular endothelial cell growth factor antibody bevacizumab [Avastin™], compounds such as those disclosed in International Patent Applications WO 97/22596, WO 97/30035, WO 97/32856 and WO 98/13354) and compounds that work by other mechanisms (for example linomide, inhibitors of integrin $\alpha v \beta 3$ function and angiostatin);
- 10 (vi) vascular damaging agents such as Combretastatin A4 and compounds disclosed in International Patent Applications WO 99/02166, WO00/40529, WO 00/41669, WO01/92224, WO02/04434 and WO02/08213;
- (vii) antisense therapies, for example those which are directed to the targets listed above, such as ISIS 2503, an anti-ras antisense;
- 15 (viii) gene therapy approaches, including for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene-directed enzyme pro-drug therapy) approaches such as those using cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi-drug resistance gene therapy; and
- 20 (ix) immunotherapy approaches, including for example ex-vivo and in-vivo approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such as interleukin 2, interleukin 4 or granulocyte-macrophage colony stimulating factor, approaches to decrease T-cell anergy, approaches using transfected immune cells such as cytokine-transfected dendritic cells, approaches using cytokine-transfected tumour cell lines
- 25 and approaches using anti-idiotypic antibodies.

Pharmacological Data

Ligand Binding Assay

- 30 [¹²⁵I]IL-8 (human, recombinant) was purchased from Amersham, U.K. with a specific activity of 2,000Ci/mmol. All other chemicals were of analytical grade. High levels of hrCXCR2 were expressed in HEK 293 cells (human embryo kidney 293 cells ECACC No. 85120602) (Lee *et al.* (1992) *J. Biol. Chem.* 267 pp16283-16291). hrCXCR2 cDNA was amplified and

-30-

- cloned from human neutrophil mRNA. The DNA was cloned into PCRScript (Stratagene) and clones were identified using DNA. The coding sequence was sub-cloned into the eukaryotic expression vector RcCMV (Invitrogen). Plasmid DNA was prepared using Quiagen Megaprep 2500 and transfected into HEK 293 cells using Lipofectamine reagent (Gibco BRL). Cells of the highest expressing clone were harvested in phosphate-buffered saline containing 0.2%(w/v) ethylenediaminetetraacetic acid (EDTA) and centrifuged (200g, 5min.). The cell pellet was resuspended in ice cold homogenisation buffer [10mM HEPES (pH 7.4), 1mM dithiothreitol, 1mM EDTA and a panel of protease inhibitors (1mM phenyl methyl sulphonyl fluoride, 2µg/ml soybean trypsin inhibitor, 3mM benzamidine, 0.5µg/ml leupeptin and 100µg/ml bacitracin)] and the cells left to swell for 10 minutes. The cell preparation was disrupted using a hand held glass mortar/PTFE pestle homogeniser and cell membranes harvested by centrifugation (45 minutes, 100,000g, 4°C). The membrane preparation was stored at -70°C in homogenisation buffer supplemented with Tyrode's salt solution (137mM NaCl, 2.7mM KCl, 0.4mM NaH₂PO₄), 0.1%(w/v) gelatin and 10%(v/v) glycerol.
- All assays were performed in a 96-well MultiScreen 0.45µm filtration plates (Millipore, U.K.). Each assay contained ~50pM [¹²⁵I]IL-8 and membranes (equivalent to ~200,000 cells) in assay buffer [Tyrode's salt solution supplemented with 10mM HEPES (pH 7.4), 1.8mM CaCl₂, 1mM MgCl₂, 0.125mg/ml bacitracin and 0.1%(w/v) gelatin]. In addition, a compound of formula (I) according to the Examples was pre-dissolved in DMSO and added to reach a final concentration of 1%(v/v) DMSO. The assay was initiated with the addition of membranes and after 1.5 hours at room temperature the membranes were harvested by filtration using a Millipore MultiScreen vacuum manifold and washed twice with assay buffer (without bacitracin). The backing plate was removed from the MultiScreen plate assembly, the filters dried at room temperature, punched out and then counted on a Cobra β -counter.
- The compounds of formula (I) according to the Examples 1 – 138 were found to have pIC₅₀ values of greater than (>) 5.0. For example, Examples 3, 4 and 116 were found to have pIC₅₀ values of 7.10, 7.10 and 6.80 respectively.

Intracellular Calcium Mobilisation Assay

- Human neutrophils were prepared from EDTA-treated peripheral blood, as previously described (Baly *et al.* (1997) *Methods in Enzymology* 287 pp70-72), in storage buffer [Tyrode's salt solution (137mM NaCl, 2.7mM KCl, 0.4mM NaH₂PO₄) supplemented with 5.7mM glucose and 10mM HEPES (pH 7.4)].

The chemokine GRO δ (human, recombinant) was purchased from R&D Systems (Abingdon, U.K.). All other chemicals were of analytical grade. Changes in intracellular free calcium were measured fluorometrically by loading neutrophils with the calcium sensitive fluorescent dye, fluo-3, as described previously (Merritt *et al.* (1990) *Biochem. J.* 269, pp513-519). Cells were loaded for 1 hour at 37°C in loading buffer (storage buffer with 0.1%(w/v) gelatin) containing 5 μ M fluo-3 AM ester, washed with loading buffer and then resuspended in Tyrode's salt solution supplemented with 5.7mM glucose, 0.1%(w/v) bovine serum albumin (BSA), 1.8mM CaCl₂ and 1mM MgCl₂. The cells were pipetted into black walled, clear bottom, 96 well micro plates (Costar, Boston, U.S.A.) and centrifuged (200g, 5 minutes, room temperature).

A compound of formula (I) according to the Examples was pre-dissolved in DMSO and added to a final concentration of 0.1%(v/v) DMSO. Assays were initiated by the addition of an A₅₀ concentration of GRO δ and the transient increase in fluo-3 fluorescence (δ_{Ex} = 490nm and δ_{Em} = 520nm) monitored using a FLIPR (Fluorometric Imaging Plate Reader, Molecular Devices, Sunnyvale, U.S.A.).

The compounds of formula (I) according to the Examples were tested and found to be antagonists of the CXCR2 receptor in human neutrophils.

The invention will now be illustrated by the following non-limiting Examples in which, unless stated otherwise:

- (i) when given Nuclear Magnetic Resonance (NMR) spectra were measured on a Varian Unity Inova 300 or 400 MHz spectrometer. ¹H NMR data is quoted in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard.
- (ii) Mass Spectrometry (MS) spectra were measured on a Finnigan Mat SSQ7000 or Micromass Platform spectrometer.
- (iii) the title and sub-titled compounds of the Examples and methods were named using the ACD/Name program (version 4.55) from Advanced Chemical Development Inc, Canada.
- (iv) Normal phase column chromatography and normal phase HPLC was conducted using a silica column. Reverse phase High Pressure Liquid Chromatography (HPLC) purification was performed using either a Waters Micromass LCZ with a Waters 600 pump controller, Waters 2487 detector and Gilson FC024 fraction

-32-

collector or a Waters Delta Prep 4000 or a Gilson Auto Purification System, using a Symmetry, NovaPak or Ex-Terra reverse phase silica column.

(v) The following abbreviations are used:

AcOH acetic acid

5 CHCl₃ chloroform

DCM dichloromethane

DMF *N,N*-dimethylformamide

DMSO dimethylsulfoxide

Et₂O diethyl ether

10 EtOAc ethyl acetate

MgSO₄ magnesium sulfate

NMP 1-methylpyrrolidin-2-one

THF tetrahydrofuran


H₂O water

15

Example 1

***N*-(2-[(2,3-difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)methanesulfonamide**

5



- Methanesulfonyl chloride (0.16ml) was added to a solution of the subtitle product of step iv) (0.40g) and *N,N*-diisopropylethylamine (0.5ml) in DCM (15ml) and stirring maintained for 2h. The reaction solution was extracted with H₂O (2 x 20ml) and the organics combined, dried (MgSO₄) and concentrated to yield a brown oil. The residue was diluted in THF (10ml) and treated with 1M tetrabutylammonium fluoride in THF (2ml) for 30min at room temperature. The volatiles were removed *in vacuo* and the residue partitioned between EtOAc (30ml) and saturated ammonium chloride solution (30ml). The aqueous layer was further extracted with EtOAc (2 x 20ml), the organics combined, dried (MgSO₄) and concentrated to yield a white solid. This material was further purified by silica gel chromatography and then reverse phase HPLC (acetonitrile / 0.02M ammonium hydroxide (90% to 5% aqueous phase) to yield the title compound as a white solid. Yield: 25mg.
- MS APCI(+ve) 405 [M+H]⁺
- ¹H NMR δ (DMSO) 7.41-7.12 (3H, m), 5.79 (1H, s), 4.70 (1H, br. s), 4.38 (2H, s), 3.41-3.25 (2H, m), 3.22 (3H, s), 1.05 (3H, d).

The intermediates for this compound were prepared as follows:

i) 6-Amino-2-[(2,3-difluorobenzyl)thio]pyrimidin-4(3*H*)-one

- An aqueous solution of potassium hydroxide (4.61g) in H₂O (25ml) was added to a suspension of 4-amino-6-hydroxy-2-mercaptopyrimidine monohydrate (11.26g) in DMF (50ml). Stirring was maintained for 30min during which time a solution was obtained, before the dropwise addition of a solution of 2,3-difluorobenzyl bromide (14.46g) in THF (10ml). After stirring for 20h the slurry was diluted with H₂O (500ml) and stirred for 30min before

-34-

filtering. The filtrate was washed with H₂O (4 x 100ml) and iso-hexane (4 x 100ml) before drying *in vacuo* for 24h to afford the subtitle compound as a white solid. Yield: 14.1g.

MS APCI(+ve) 309 [M+CH₃COO]⁺

5 **ii) 6-Chloro-2-[(2,3-difluorobenzyl)thio]pyrimidin-4-amine**

N,N-Dimethylaniline (5ml) was added to a solution of the subtitle product of step i) in phosphorus oxychloride (50ml) and heated at reflux for 2h. The reaction was allowed to cool before pouring into hot H₂O (500ml) and stirring the mixture for 2h. This mixture was extracted with DCM (3 x 250ml) and the organics combined, dried (MgSO₄) and concentrated

10 *in vacuo* to afford the subtitle compound as a green foam. This crude product was used directly in step iii). Yield: 12.3g.

MS: APCI(+ve) 329 [M+CH₃COO]⁺

iii) (2*R*)-2-({6-Amino-2-[(2,3-difluorobenzyl)thio]pyrimidin-4-yl}amino)propan-1-ol

15 *N,N*-Diisopropylethylamine (1.92ml) was added to a solution of *R*-alaninol (2.0ml) and the subtitle product of step ii) (1.9g) in NMP (10ml) and stirred at 100°C for five days before pouring into H₂O (200ml) and filtration of the precipitate. This solid was dried *in vacuo* to afford the subtitle compound as a yellow solid. Yield: 1.80g.

MS: APCI(+ve) 327 [M+H]⁺

20

iv) *N*-((1*R*)-2-({*tert*-Butyl(dimethyl)silyl}oxy)-1-methylethyl)-2-[(2,3-difluorobenzyl)thio]pyrimidine-4,6-diamine


Imidazole (1.2g) was added to a solution of *tert*-butyldimethylsilyl chloride (2.83g) and the subtitle product of step iii) (1.8g) in DMF (10ml). The reaction was stirred for 20h before
25 partitioning between EtOAc (100ml) and H₂O (200ml). The aqueous was extracted further with EtOAc (2 x 100ml), the organics combined, washed with H₂O (100ml), brine (100ml), dried (MgSO₄) and concentrated *in vacuo* to a crude solid. This material was purified by column chromatography (50% Et₂O / iso-hexane) to afford the subtitle compound as a yellow oil. Yield: 1.80g.

30 MS: APCI(+ve) 441 [M+H]⁺

-35-

Example 2

***N*-(2-[(2,3-Difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-1-methyl-1*H*-imidazole-4-sulfonamide**



5

1-Methyl-1*H*-imidazole-4-sulfonyl chloride was added to a solution of the subtitle product of Example 1 step iv) (0.40g) and 4-dimethylaminopyridine (0.12g) in pyridine (10ml) at room temperature and stirred for 20h. The reaction mixture was partitioned between DCM (50ml) and copper (II) sulfate solution (60ml). The aqueous was extracted further with DCM, the organics combined, dried (MgSO₄) and concentrated *in vacuo*. The resulting oil was diluted in THF (10ml) and treated with tetrabutylammonium fluoride (1M in THF, 2ml) for 30min at room temperature. The volatiles were removed *in vacuo* and the residue partitioned between EtOAc (20ml) and saturated ammonium chloride solution (20ml). The aqueous was further extracted with EtOAc (2 x 20ml), the organics combined, dried (MgSO₄) and concentrated to yield a crude white solid. This material was further purified by reverse phase HPLC (acetonitrile / 0.02M ammonium hydroxide (90% to 5% aqueous phase)) to yield the title compound as a white solid. Yield: 60mg.


MS APCI(+ve) 471 [M+H]⁺
¹H NMR δ (DMSO) 7.83 (m, 1H), 7.75 (s, 1H), 7.33 (m, 3H), 7.11 (m, 2H), 5.92 (s, 1H), 4.69 (s, 1H), 4.32 (s, 2H), 3.96 (s, 1H), 3.66 (s, 3H), 3.40 - 3.20 (m, 2H), 1.03 (d, 3H)

Example 3

***N*-(2-(Benzylthio)-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-methanesulfonamide**

25

-36-



A solution of the subtitle product of step iii) (0.18g) in THF (10ml) was treated with tetrabutylammonium fluoride (1M in THF, 2ml) for 2h at room temperature. The volatiles were removed *in vacuo* and the residue partitioned between EtOAc (20ml) and saturated ammonium chloride solution (20ml). The aqueous was further extracted with EtOAc (2 x 20ml), the organics combined, dried (MgSO₄) and concentrated to yield a crude white solid. This material was further purified by reverse phase HPLC (acetonitrile / 0.02M ammonium hydroxide (90% to 5% aqueous phase)) to yield the title compound as a white solid. Yield: 25mg.

MS APCI(+ve) 369 [M+H]⁺

¹H NMR δ_(DMSO) 7.41 (d, 2H), 7.30 (t, 2H), 7.23 (t, 2H), 5.78 (s, 1H), 4.71 (t, 1H), 4.32 (s, 2H), 3.40 (dt, 1H), 3.29 (m, 1H), 3.18 (s, 3H), 1.07 (d, 3H).

The intermediates for this compound were prepared as follows:

i) (2R)-2-[[6-Amino-2-(benzylthio)pyrimidin-4-yl]amino]propan-1-ol

N,N-Diisopropylethylamine (6.0ml) was added to a solution of *R*-alaninol (12.0ml) and 2-(benzylthio)-6-chloropyrimidin-4-amine (1.9g) (Nugent, R.A., *et al.* PCT Int. Appl. 1996. 252pp. WO9635678-A1) in NMP (6ml) and stirred at 100°C for three days before pouring into H₂O (200ml) and filtration of the precipitate. This solid was dried *in vacuo* to afford the subtitle compound as a pale sandy yellow solid. Yield: 4.1g.

MS: APCI(+ve) 291 [M+H]⁺

ii) 2-(Benzylthio)-*N*-((1*R*)-2-[[*tert*-butyl(dimethyl)silyl]oxy]-1-methylethyl)pyrimidine-4,6-diamine

Imidazole (0.29g) was added to a solution of *tert*-butyldimethylsilyl chloride (0.34g) and the subtitle product of step i) (0.6g) in DMF (10ml). The reaction was stirred for 24h before addition of a further equivalent of *tert*-butyldimethylsilyl chloride and imidazole. After stirring for an additional 24h the reaction mixture was partitioned between EtOAc (100ml)

-37-

and H₂O (200ml). The aqueous was extracted further with EtOAc (3 x 100ml), the organics combined, washed with H₂O (100ml), brine (100ml), dried (MgSO₄) and concentrated to a crude solid. This material was purified by column chromatography (1:1 Et₂O / iso-hexane) to afford the subtitle compound as a yellow oil. Yield: 0.50g.

5 MS: APCI(+ve) 405 [M+H]⁺

iii) *N*-{2-(Benzylthio)-6-[[((1*R*)-2-[[*tert*-butyl(dimethyl)silyl]oxy}-1-methylethyl)amino]-pyrimidin-4-yl]methanesulfonamide


Methanesulfonyl chloride (85 μl) was added to a solution of the subtitle product of step ii)

10 (0.20g) and *N,N*-diisopropylethylamine (0.26ml) in DCM (10ml) at 0°C. The ice-bath was removed and stirring maintained for 2h. The reaction solution was extracted with H₂O (2 x 20ml) and the organics dried (MgSO₄) and concentrated to yield a brown oil. The residue was diluted in methanol (10ml) and treated with potassium carbonate (0.15g) for 2h at room temperature. The volatiles were removed *in vacuo* and the residue partitioned between
15 EtOAc (20ml) and H₂O (20ml). The aqueous was further extracted with EtOAc (2 x 20ml), the organics combined, dried (MgSO₄) and concentrated to yield the subtitle compound as a crude white solid. This material was used directly in the next step. Yield: 0.23g.

MS APCI(+ve) 483 [M+H]⁺

20 Example 4

***N*-(2-(Benzylthio)-6-[[((1*R*)-2-hydroxy-1-methylethyl)amino}pyrimidin-4-yl)benzenesulfonamide**



25

Phenylsulfonyl chloride (0.20g) was added to a solution of the subtitle product of Example 3 step ii) (0.40g) and 4-dimethylamino pyridine (0.17g) in pyridine (10ml) was stirred for 24h at room temperature. The reaction was quenched with 10% potassium carbonate solution (10ml) and the aqueous extracted with EtOAc (2 x 20ml). The crude material was dissolved

-38-

in THF (10ml) and treated with tetrabutylammonium fluoride (1M in THF, 5ml) for 15min at room temperature. The reaction was quenched with 1M hydrochloric acid (10ml) and the aqueous extracted with EtOAc (2 x 20ml). The organics were then combined, washed with brine (50ml), dried (MgSO₄) and concentrated to yield a crude gum which was purified by column chromatography (2% methanol / DCM) to afford a gum. This material was treated with ethanol (25ml) and H₂O (5ml) and the volatiles removed under reduced pressure to yield the title compound as a white solid. Yield: 0.39g.


MS APCI(+ve) 431 [M+H]⁺

¹H NMR δ (DMSO) 7.87 (d, 2H), 7.60 (m, 3H), 7.35 (d, 2H), 7.28 (t, 2H), 7.22 (m, 1H), 5.89 (s, 1H), 4.70 (s, 1H), 4.21 (s, 2H), 4.01 (s, 1H), 3.40 - 3.21 (m, 2H), 1.04 (d, 3H).

Example 5

N-(2-(Benzylthio)-6-{[(1*R*)-1-methylpropyl]amino}pyrimidin-4-yl)-3,5-dimethylisoxazole-4-sulfonamide

15




3,5-Dimethylisoxazole-4-sulfonyl chloride (0.60g) was added to a solution of the subtitle product of Example 3 step ii) (50mg) in pyridine (0.5ml) and *N,N*-dimethylamino-pyridine (16mg). The reaction mixture was stirred overnight. To this reaction was added excess aqueous hydrochloric acid (1M) and stirred for 1h. The solvent was evaporated and the residue diluted in EtOAc (100ml). This was washed with H₂O (2 x 20ml) and brine (10ml). The organic layer was dried (MgSO₄) and concentrated to yield a white solid. This material was purified by reverse phase HPLC (gradient 90% to 5% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a solid. Yield: 24mg.

MS APCI(+ve) 450 [M+H]⁺

¹H NMR δ (DMSO) 7.26-7.30 (5H, m), 5.73 (1H, s), 5.30 (1H, bs), 4.30 (2H, s), 4.0 (1H, br. s), 3.66 (1H, m), 3.54-3.59 (1H, m), 2.65 (3H, s), 2.39 (3H, s), 1.21 (3H, d).

Example 6***N*-(2-(Benzythio)-6-{[(1*R*)-1-methylpropyl]amino}pyrimidin-4-yl)-1-phenylmethanesulfonamide**


5



Phenylmethanesulfonyl chloride (0.28g) was added to a solution of the subtitle product of Example 3 step ii) (0.1g) in pyridine (1ml) and *N,N*-dimethylaminopyridine (30mg). The reaction mixture was stirred for 6h. To the reaction mixture was added tetrabutylammonium fluoride (7ml, 1M in THF) and stirred for 16h. The solvent was evaporated and the residue diluted in EtOAc (100ml) and hydrochloric acid (1M). This was washed with H₂O (2 x 20ml) and brine (10ml). The organics were dried (MgSO₄) and concentrated to yield a solid. This material was purified by reverse phase HPLC (90% to 5% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a solid. Yield: 30mg.

MS APCI(+ve) 445 [M+H]⁺

¹H NMR δ (CDCl₃) 7.24-7.40 (10H, m), 5.76 (1H, s), 5.02 (1H, bs), 4.43 (2H, s), 4.31 (2H, s), 4.02 (1H, br. s), 3.65-3.70 (1H, m), 3.51-3.56 (1H, m), 1.21 (3H, d).

20 **Example 7*****N*-(2-(2-Chlorobenzythio)-6-{[(1*R*)-1-methylpropyl]amino}pyrimidin-4-yl)-2-chlorobenzenesulfonamide**


The title compound was synthesised according to the procedure of Example 6 using 2-chlorobenzenesulfonyl chloride (0.31g), the subtitle product of Example 3 step ii) (0.1g), *N,N*-dimethylaminopyridine (30mg) and tetrabutylammonium fluoride (4ml, 1M in THF) to yield
5 the title compound as a white solid. Yield: 30mg.

MS APCI(+ve) 465 $[M+H]^+$

^1H NMR $\delta_{(\text{CDCl}_3)}$ 8.14 (1H, d), 7.48 (2H, bs), 7.22-7.35 (6H, m), 5.84 (1H, s), 5.03 (1H, bs), 4.25 (2H, s), 3.98 (1H, bs) 3.54-3.66 (1H, m), 3.49-3.54 (1H, m), 1.21 (3H, d).

10 Example 8

***N*-(2-(Benzylthio)-6-{[(1*R*)-1-methylpropyl]amino}pyrimidin-4-yl)-4-cyanobenzenesulfonamide**



15

The title compound was synthesised according to the procedure of Example 6 using 4-cyanobenzenesulfonyl chloride (0.30g), the subtitle product of Example 3 step ii) (0.1g), *N,N*-dimethylaminopyridine (30mg) and tetrabutylammonium fluoride (4ml, 1M in THF) to yield the title compound as a white solid. Yield: 45mg.


20 MS APCI(+ve) 456 $[M+H]^+$

^1H NMR $\delta_{(\text{CDCl}_3)}$ 8.05 (2H, d), 7.74 (2H, d), 7.19-7.34 (5H, m), 5.85 (1H, s), 5.44 (1H, d), 4.13 (2H, s), 4.00 (1H, bs), 3.68-3.73 (1H, m), 3.54-3.59 (1H, m), 1.21 (3H, d).

Example 9

25 ***N*-(2-(Benzylthio)-6-{[(1*R*)-1-methylpropyl]amino}pyrimidin-4-yl)-4-chlorobenzenesulfonamide**

-41-



The title compound was synthesised according to the procedure of Example 6 using 4-chlorobenzenesulfonyl chloride (0.31g), the subtitle product of Example 3 step ii) (0.1g), *N,N*-dimethylaminopyridine (30mg) and tetrabutylammonium fluoride (4ml, 1M in THF) to yield the title compound as a white solid. Yield: 18mg.


MS APCI(+ve) 465 [M+H]⁺

¹H NMR δ (CDCl₃) 7.80 (2H, d), 7.40 (2H, d), 7.20-7.40 (5H, m), 5.90 (1H, s), 5.15 (1H, bs), 4.20 (2H, s), 4.00 (1H, bs) 3.60-3.80 (1H, m), 3.42-3.60 (1H, m), 1.21 (3H, d).

10

Example 10

***N*-(2-(Benzylthio)-6-{[(1*R*)-1-methylpropyl]amino}pyrimidin-4-yl)-2-cyanobenzenesulfonamide**



15

2-Cyanobenzenesulfonyl chloride (0.30g) was added to the solution of the subtitle product of Example 3 step ii) (0.1g) in pyridine (1ml) and *N,N*-dimethylaminopyridine (30mg). The reaction mixture was stirred for 4h. To the reaction mixture was added tetrabutylammonium fluoride (4ml, 1M in THF) and stirred for 16h. To this reaction was added aqueous hydrochloric acid (1M, 30ml). The solvent was evaporated and the residue diluted in EtOAc (100ml). This was washed with H₂O (3 x 20ml) and brine (10ml). The organic layer was dried (MgSO₄) and concentrated to yield a solid. The residue was purified by column

-42-

chromatography (EtOAc / iso-hexane 7:3) followed by column chromatography (EtOAc) to yield the title compound. Yield: 30mg.


MS APCI(+ve) 456 [M+H]⁺

¹H NMR δ (CDCl₃) 8.20 (1H, d), 7.58-7.62 (1H, m), 7.62-7.78 (1H, m), 7.78-8.00 (1H, m), 7.19-7.35 (5H, m), 5.80 (1H, s), 5.48 (1H, bs), 4.25 (2H, s), 3.50-3.80 (2H, m), 1.21 (3H, d).

Example 11

***N*-(2-(Benzylthio)-6-{[(1*R*)-1-methylpropyl]amino}pyrimidin-4-yl)-3-cyanobenzenesulfonamide**

10



3-Cyanobenzenesulfonyl chloride (0.30g) was added to the solution of the subtitle product of Example 3 step ii) (0.1g) in pyridine (1ml) and *N,N*-dimethylaminopyridine (30mg). The reaction mixture was stirred for 4h. To the reaction mixture was added tetrabutylammonium fluoride (4ml, 1M in THF) and stirred for 16h. To this reaction was added aqueous hydrochloric acid (1M, 30ml). The solvent was evaporated and the residue diluted in EtOAc (100ml). This was washed with H₂O (3 x 20ml) and brine (10ml). The organic layer was dried (MgSO₄) and concentrated to yield a solid. The residue was purified by column chromatography (EtOAc / iso-hexane 7:3 to 1:1) to yield the title compound as a white solid. Yield: 45mg.


MS APCI(+ve) 456 [M+H]⁺

¹H NMR δ (CDCl₃) 8.20 (1H, s), 8.15 (1H, d), 7.80 (1H, d), 7.60 (1H, t), 7.20-7.40 (6H, m), 5.90 (1H, s), 5.50 (1H, d), 4.32 (2H, s), 3.50-3.80 (2H, m), 1.21 (3H, d).

25

Example 12

***N*-(2-(Benzylthio)-6-{[(1*R*)-1-methylpropyl]amino}pyrimidin-4-yl)-3-chlorobenzenesulfonamide**



3-Chlorobenzenesulfonyl chloride (0.30g) was added to the solution of the subtitle product of Example 3 step ii) (0.1g) in pyridine (1ml) and *N,N*-dimethylaminopyridine (30mg). The reaction mixture was stirred for 4h. To the reaction mixture was added tetrabutylammonium fluoride (4ml, 1M in THF) and stirred for 16h. To this reaction was added aqueous hydrochloric acid (1M, 30ml). The solvent was evaporated and the residue diluted in EtOAc (100ml). This was washed with H₂O (3 x 20ml) and brine (10ml). The organic layer was dried (MgSO₄) and concentrated to yield a solid. This material was purified by reverse phase HPLC (90% to 5% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a solid. Yield: 30mg.


MS APCI(+ve) 465 [M+H]⁺

¹H NMR δ (CDCl₃) 7.93 (1H, m), 7.80 (1H, d), 7.55 (1H, m), 7.45 (1H, t), 7.20-7.40 (5H, m), 5.90 (1H, s), 5.20 (1H, bd), 4.32 (2H, s), 4.00 (1H, bs), 3.60-3.70 (1H, m), 3.45-3.60 (1H, m), 1.21 (3H, d).

Example 13

N-(5-(((2-(Benzylthio)-6-(((1*R*)-1-methylpropyl)amino}pyrimidin-4-yl)amino)sulfonyl)-4-methyl-1,3-thiazol-2-yl)acetamide

20




-44-

- 2-(Acetylamino)-1,3-thiazole-4-sulfonyl chloride (0.19g) was added to a solution of the subtitle product of Example 3 step ii) (0.2g) in pyridine (4ml) and *N,N*-dimethylaminopyridine (60mg). The reaction mixture was stirred for 4 days. To this reaction more sulfonyl chloride (0.75g) was added and stirred for 2 days. To this reaction was added aqueous hydrochloric acid (1M, 20ml) and THF (20ml). This was stirred for 18h before the solvent was evaporated and the residue diluted in EtOAc (100ml). This was washed with H₂O (3 x 20ml) and brine (10ml). The organic layer was dried (MgSO₄) and concentrated to yield a solid. This material was purified by reverse phase HPLC (90% to 5% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a solid. Yield: 20mg.
- 10 MS APCI(+ve) 509 [M+H]⁺
¹H NMR δ (DMSO) 7.20-7.37 (5H, m), 5.75 (1H, bs), 4.69 (1H, bs), 4.27 (2H, bs), 3.31-3.39 (2H, m), 2.50 (3H, s), 2.12 (3H, s), 1.21 (3H, d).

Example 14

- 15 *N*-(2-(Benzylthio)-6-{[(1*R*)-1-methylpropyl]amino}pyrimidin-4-yl)-2-(methylsulfonyl)benzenesulfonamide



- 20 2-(Methylsulfonyl)benzenesulfonyl chloride (0.19g) was added to a solution of the subtitle product of Example 3 step ii) (0.2g) in pyridine (4ml) and *N,N*-dimethylaminopyridine (59mg). The reaction mixture was stirred for 4 days at room temperature. The solvent was removed, THF (2ml) and sodium hydroxide (10%, 3ml) added and stirring maintained for 2h. The volatiles were removed *in vacuo* and the aqueous residue extracted with EtOAc (2 x 20ml) and evaporated. To this residue was added aqueous hydrochloric acid (1M, 30ml) and THF (10ml). This was stirred at room temperature for 1h. The mixture was extracted with EtOAc (2 x 30ml). This was washed with H₂O (2 x 20ml) and brine (20ml). The organic layer was dried (MgSO₄) and concentrated to yield a solid. This material was purified by

-45-

reverse phase HPLC (90% to 5% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a solid. Yield: 30mg.


MS APCI(+ve) 510 $[M+H]^+$

^1H NMR δ (CDCl_3) 8.31-8.34 (1H, dd), 8.23-8.26 (1H, dd), 7.30-7.40 (2H, d), 7.18-7.30 (3H, m),
5 6.18 (1H, s), 4.90 (1H, d), 4.20 (2H, s), 4.00 (1H, bs), 3.61-3.65 (1H, m), 3.45-3.60 (1H, m),
3.45 (3H, s), 1.21 (3H, d).

Example 15

N-(2-(Benzylthio)-6-{[(1*R*)-1-methylpropyl]amino}pyrimidin-4-yl)-4-

10 (methylsulfonyl)benzenesulfonamide



The title compound was synthesised according to the procedure of Example 14
15 using 4-methylsulfonylbenzenesulfonylchloride (0.19g), the subtitle product of Example 3
step ii) (0.2g) and *N,N*-dimethylaminopyridine (60mg) to yield the title compound as a white
solid. Yield: 60mg.


MS APCI(+ve) 510 $[M+H]^+$

^1H NMR δ (CDCl_3) 8.09-8.12 (1H, d), 7.97-8.00 (2H, d), 7.20-7.30 (5H, m), 5.90 (1H, s), 5.65
20 (1H, bs), 4.28 (2H, s), 4.00 (1H, bs), 3.69-3.71 (1H, m), 3.49-3.60 (1H, m), 3.05 (3H, s), 1.17-
1.19 (3H, d).

Example 16

N-(2-(Benzylthio)-6-{[(1*R*)-1-methylpropyl]amino}pyrimidin-4-yl)propane-1-

25 sulfonamide



Propane-1-sulfonyl chloride (0.14g) in DCM (1ml) was added to a solution of the subtitle product of Example 3 step ii) (0.2g) in DCM (3ml) and *N,N*-diisopropylethylamine (0.14g) at 5 0°C. The reaction mixture was stirred for 24h at room temperature. To the reaction mixture more *N,N*-diisopropylethylamine (0.14g) was added and the reaction mixture was stirred for an additional 24h. The DCM was removed under reduced pressure and the residue dissolved in THF (2ml). To this mixture sodium hydroxide (10%, 2ml) was added and stirred overnight. The reaction mixture was diluted with EtOAc (50ml). The organic layer was 10 separated from the aqueous layer and evaporated to dryness. To the residue was added aqueous hydrochloric acid (1M, 30ml) and THF (10ml). This was stirred at room temperature for 3h. The mixture was extracted with EtOAc (2 x 50ml). The organics were washed with brine (30ml). The combined organic layers were dried (MgSO₄) and concentrated. This material was purified by reverse phase HPLC (90% to 5% 0.02M ammonium hydroxide / 15 acetonitrile) to yield the title compound as a white solid. Yield: 30mg.

MS APCI(+ve) 397 [M+H]⁺


¹H NMR δ (CDCl₃) 7.21-7.41 (5H, m), 5.94 (1H, s), 5.05 (1H, d), 4.30 (2H, s), 4.10 (1H, bs), 3.68-3.73 (1H, m), 3.53-3.60 (1H, m), 3.16-3.21 (2H, t), 1.77-1.90 (2H, m), 1.21 (3H, d), 1.00 (3H, t).

20

Example 17

***N*-(2-(Benzylthio)-6-{[(1*R*)-1-methylpropyl]amino}pyrimidin-4-yl)-5-chloro-1,3-dimethyl-1*H*-pyrazole-4-sulfonamide**

-47-




3-Chloro-1,5-dimethyl-1*H*-pyrazole-4-sulfonyl chloride (0.34g) was added to the subtitle product of Example 3 step ii) (0.1g) in pyridine (1ml) and *N,N*-dimethylaminopyridine (30mg). The reaction mixture was stirred overnight at room temperature. To this reaction was added tetrabutylammonium fluoride (4ml, 1M in THF) and stirred for 3h. To this mixture was added aqueous hydrochloric acid (30ml, 1M) and extracted with EtOAc (2 x 30ml) then brine (20ml). The combined organic layer was dried (MgSO₄) and concentrated. This material was purified by column chromatography (EtOAc / iso-hexane (1:1) to EtOAc) to afford the title compound as a white solid. Yield: 30mg.

MS APCI(+ve) 484[M+H]⁺

¹H NMR δ (CDCl₃) 7.22-7.37 (5H, m), 5.87 (1H, s), 4.95 (1H, br,s), 4.27 (2H, m), 4.0 (1H, br, s), 3.80 (3H, s), 3.66-3.70 (1H, m), 3.52-3.56 (1H, m), 2.44 (3H, s), 1.20 (3H, d).

15 **Example 18**

N-(2-(Benzylthio)-6-((1*R*)-1-methylpropylamino)pyrimidin-4-yl)-1,3,5-trimethyl-1*H*-pyrazole-4-sulfonamide



20

The title compound was synthesised according to the procedure of Example 17 using 1,3,5-trimethyl-1*H*-pyrazole-4-sulfonyl chloride (0.30g), the subtitle product of

-48-


Example 3 step ii) (0.1g), *N,N*-dimethylaminopyridine (30mg) and tetrabutylammonium fluoride (4ml, 1M in THF) to yield the title compound as a white solid. Yield: 20mg.

MS APCI(+ve) 463[M+H]⁺

¹H NMR δ (CDCl₃) 7.21-7.36 (5H, m), 5.79 (1H, s), 4.95 (1H, d), 4.27 (2H, s), 4.0 (1H, br. s),
5 3.70 (3H, s), 3.64-3.68 (1H, m), 3.50-3.54 (1H, m), 2.41 (3H, s), 2.39 (3H, s), 1.20 (3H, d).

Example 19

N-{2-(Benzylthio)-6-[(2-hydroxyethyl)amino]pyrimidin-4-yl}methanesulfonamide



10

To the subtitle product of step iii) (0.20g) was added ethanolamine (3.0ml) and the reaction was heated at 100°C for 1h. To the reaction was added EtOAc (50ml) and H₂O (50ml). The organic layer was separated and washed with H₂O (2 x 20ml) and brine (20ml).

15 The organic layer was dried (MgSO₄), the solids were filtered and the solvent removed under reduced pressure to give a solid. This was purified by column chromatography (EtOAc / iso-hexane 1:1) to yield the title compound as a solid. Yield: 30mg.

MS APCI(+ve) 355[M+H]⁺

¹H NMR δ (CDCl₃) 7.19-7.40 (5H, m), 5.91 (1H, s), 5.45 (1H, t), 4.33 (2H, s), 3.77 (2H, t), 3.52
20 (2H, m), 3.13(3H, s).

The intermediates for this compound were prepared as follows:

i) 2-(Benzylthio)pyrimidine-4,6-diol

A solution of sodium hydroxide (3.30g) in ethanol / H₂O (60ml / 60ml) was added to 2-
25 mercaptopyrimidine-4,6-diol (10.00g) and the mixture stirred for 10min. Benzyl bromide (13.45g) was then added dropwise and the mixture heated at 60°C for 2h. The reaction was cooled to 0°C for 1h before the precipitate was filtered and washed with H₂O (100ml) and then dried *in vacuo* to afford the subtitle compound as a cream solid. Yield: 15.0g.

¹H NMR δ (DMSO) 7.41-7.46 (2H, m), 7.20-7.40 (4H, m), 4.39 (2H, s).

ii) 2-(Benzylthio)-4,6-dichloropyrimidine

N,N-Dimethylaniline (7ml) was added to the slurry of the subtitle product of step i) (5.0g) in phosphorus oxychloride (35ml) and heated at reflux for 10h. The reaction was allowed to cool and excess phosphorus oxychloride was removed *in vacuo* before pouring onto ice. This mixture was extracted with EtOAc (200ml) and washed with brine (2x100ml), dried (MgSO₄) and concentrated *in vacuo* to afford the crude product. This crude product was purified by column chromatography (EtOAc / iso-hexane (10 to 20%)) to yield the title compound. Yield: 4.10g.

¹H NMR δ (CDCl₃) 7.40-7.42 (2H, m), 7.20-7.30 (4H, m), 4.38 (2H, s).

iii) *N*-[2-(Benzylthio)-6-chloropyrimidin-4-yl]methanesulfonamide


To methane sulphonamide (1.47g) in DMF (30ml) was added 60% sodium hydride (0.59g) at room temperature and stirred for 1h. To this mixture was then added the subtitle product of step ii) and the mixture stirred at room temperature for 6h. To the reaction mixture was added EtOAc (50ml) and aqueous hydrochloric acid (50ml, 1M). The organic layer was separated and washed with brine (20ml), combined, dried (MgSO₄) and evaporated to dryness under reduced pressure. The residue was purified by column chromatography (20 to 50% EtOAc / iso-hexane) to afford the subtitle compound as a yellow oil. Yield: 1.60g.

MS APCI (+ve) 330 [M+H]⁺

¹H NMR δ (DMSO) 7.40-7.42 (2H, d), 7.20-7.40 (3H, m), 6.70 (1H, s), 4.38 (2H, s), 3.30 (3H, s).

Example 20

***N*-(2-(Benzylthio)-6-[(1*R*)-1-(hydroxymethyl)propyl]amino}pyrimidin-4-yl)methanesulfonamide**



-50-

To the subtitle product of Example 19 step iii) (0.20g) in NMP (1ml) was added (2R)-2-aminobutan-1-ol (1.0g) and the reaction was heated at 100°C for 2 days. To the reaction was added EtOAc (50ml) and H₂O (50ml). The organic layer was separated and washed with H₂O (2 x 20ml) and brine (20ml). The organic layer was combined, dried

5 (MgSO₄) and the solvent removed under reduced pressure to afford a solid. This was purified by column chromatography (EtOAc / iso-hexane 1:1) to give the title compound as a solid.


Yield: 15mg.

MS APCI(+ve) 382 [M+H]⁺

¹H NMR δ_(DMSO) 7.2-7.40 (2H, m), 7.20-7.30 (3H, m), 5.80 (1H, bs), 4.80 (1H, bs), 4.30 (2H, s), 3.85 (1H, bs), 3.30-3.45 (2H, m), 3.20 (3H, s), 1.41-1.64 (1H, m), 1.3-1.42 (1H, m), 0.83 (3H, t).

Example 21

N-(2-(Benzylthio)-6-{[2-hydroxy-1-(hydroxymethyl)ethyl]amino}-pyrimidin-4-yl)methanesulfonamide




To a solution of the subtitle product of Example 19 step iii) (0.20g) in NMP (0.5ml) was added 2-aminopropane-1,3-diol (1.0g) and the reaction was heated at 100°C for 2 days. To the reaction was added EtOAc (50ml) and H₂O (50ml). The organic layer was separated and washed with H₂O (2 x 20ml) and brine (20ml). The organic layer was dried (MgSO₄) and the solvent removed under reduced pressure to give a solid. This was purified by column chromatography (1:1 EtOAc / iso-hexane then EtOAc) to give the title compound as a solid.

25 Yield: 20mg.

MS APCI(+ve) 385 [M+H]⁺

¹H NMR δ_(DMSO) 7.40-7.42 (2H, m), 7.20-7.30 (3H, m), 5.85 (1H, s), 4.63 (1H, s), 4.32 (2H, s), 3.40 (4H, bs), 3.20 (3H, s).


Example 22***N*-(2-(Benzylthio)-6-{[(2*R*)-2-hydroxypropyl]amino}pyrimidin-4-yl)methanesulfonamide**

5

To the subtitle product of Example 19 step iii) (0.20g) in NMP (2ml) was added (2*R*)-1-amino-2-propanol (0.46g) and the reaction was heated at 80°C for 6h. To the reaction was added EtOAc (50ml) and H₂O (20ml). This solution was acidified with aqueous hydrochloric acid. The organic layer separated and washed with H₂O (2x20ml), brine (20ml) and the organic layer was dried (MgSO₄) and the solvent removed under reduced pressure to give a solid. This material was purified by reverse phase HPLC (90% to 5% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a solid. Yield: 20mg.

MS APCI(+ve) 370[M+H]⁺

15 ¹H NMR δ (CDCl₃) 7.2-7.47 (5H, m), 5.90 (1H, s), 5.40 (1H, bs), 4.32 (2H, s), 3.90-4.10 (1H, m), 3.40-3.50 (1H, m), 3.21 (1H, m), 3.10 (3H, bs), 1.20 (3H, d).

Example 23***N'*-(2-(Benzylthio)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-*N,N*-dimethylsulfamide**

To the subtitle product of step i) (0.20g) was added (*R*)-alaninol (2.0ml) and the mixture heated at 80°C for 2 days. The reaction mixture was diluted in EtOAc (50ml) and

25

-52-

acidified with aqueous hydrochloric acid (1M, 20ml). The aqueous was extracted further with EtOAc (50ml). The combined organic layers were washed with H₂O (3 x 20ml) and brine (20ml). The organic layer separated and further washed with H₂O (2x20ml), brine (20ml) and the organic layer was dried (MgSO₄) and the solvent removed under reduced pressure to give a solid. This material was purified by reverse phase HPLC (90% to 5% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a solid. Yield: 20mg.

MS APCI(+ve) 398[M+H]⁺

¹H NMR δ (CDCl₃) 7.2-7.40 (5H, m), 5.92 (1H, s), 4.98 (1H, bd), 4.32 (2H, s), 4.07 (1H, bs), 3.68-3.75 (1H, m), 3.50-3.60 (1H, m), 2.87 (6H, s), 1.20 (3H, d).

10

The intermediates for the title compound were prepared as follows:

i) *N'*-[2-(Benzylthio)-6-chloropyrimidin-4-yl]-*N,N*-dimethylsulfamide

To *N,N*-dimethylsulphonamide (1.0g) in DMF (10ml) was added 60% sodium hydride (0.22g) at room temperature and the reaction heated to 50°C for 1h. This mixture was then allowed to cool to room temperature and the subtitle product of Example 19 step ii) was added in DMF (1ml). To the reaction mixture was added EtOAc (50ml) and aqueous hydrochloric acid (50ml, 1M). The organic layer was separated and washed with brine (20ml). The organic layer was dried (MgSO₄) and evaporated to dryness under reduced pressure. The residue was purified by column chromatography (EtOAc / iso-hexane 1:4) to afford the title compound as a yellow gum. Yield: 0.48g.


20

MS APCI (+ve) 359 [M+H]⁺

¹H NMR δ (CDCl₃) 7.40-7.42 (2H, m), 7.20-7.40 (3H, m), 6.80 (1H, s), 4.38 (3H, s), 2.92 (6H, s).

25 Example 24

***N*-(2-(Benzylthio)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-1-methyl-2-oxoindoline-6-sulfonamide**



1-Methyl-2-oxoindoline-5-sulfonyl chloride (0.74g) was added to the cooled solution of the subtitle product of Example 3 step ii) (0.25g) in pyridine (5ml) and *N,N*-


5 dimethylaminopyridine (75mg). The reaction mixture was stirred for 3 days at room temperature. To this mixture was added aqueous hydrochloric acid (30ml, 1M) and extracted with EtOAc (2x30ml), brine (20ml). The organic layer was dried (MgSO_4) and concentrated. This material was purified by reverse phase HPLC (90% to 5% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a solid. Yield: 10mg.

10 MS APCI(+ve) 500[M+H]⁺

¹H NMR δ (DMSO) 7.80 (1H, d), 7.70 (1H, s), 7.19-7.32 (5H, m), 7.06 (1H, d), 5.82 (1H, s), 4.65 (1H, t), 4.20 (2H, s), 3.95 (1H, bs), 3.57 (2H, s), 3.20-3.41 (2H, m), 3.10 (3H, s), 1.21 (3H, d).

15 **Example 25**

1-{[(2-(benzylthio)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)amino]sulfonyl}-*N,N*-dimethyl-*L*-prolinamide



20 To the subtitle product of step iv) (0.50g) was added (*R*)-alaninol (2 ml) and the mixture heated at 80-90°C for 3 days. The reaction mixture was taken in EtOAc (1L) and acidified with aqueous hydrochloric acid (1M, 20ml). The aqueous was evaporated to give a residue which was purified by column chromatography (2% methanol / EtOAc) to give the title

-54-

compound which was further purified by reverse phase HPLC (90% to 5% 0.02M ammonium hydroxide / acetonitrile) to afford the title compound as a white solid. Yield: 20mg.

MS APCI(+ve) 495[M+H]⁺

¹H NMR δ (DMSO) 7.39-7.42 (2H, m), 7.19-7.32 (3H, m), 5.70 (1H, s), 5.16 (1H, bs), 4.70 (1H, t), 4.30 (2H, m), 4.32 (2H, s), 3.38-3.45 (1H, m), 3.20-3.38 (3H, m), 3.04 (3H, s), 2.80 (3H, s), 1.95-2.10 (1H, m), 1.80-1.92 (1H, m), 1.62-1.80 (2H, m), 1.07 (3H, d).

The intermediates for this compound were prepared as follows:

i) 1-(tert-Butoxycarbonyl)-N,N-dimethyl-L-prolinamide

10 To 1-(tert-butoxycarbonyl)-L-proline (5.0g) in DCM (50ml) at 5°C was added dicyclohexylcarbodiimide (5.22g) and N-hydroxysuccinimide (2.91g). The mixture was stirred at this temperature for 16h. The solid was filtered and the filtrate cooled to 5°C. To this mixture was added triethylamine (9.80ml) and dimethylamine hydrochloride (2.80g). The mixture was stirred at room temperature for 2 days. H₂O (50ml) was added and the
15 phases were separated and the organics were washed with saturated sodium carbonate (2x20ml) and brine (20ml). This was then dried (MgSO₄) and evaporated to dryness to afford the title compound as a white solid. Yield: 6.0g.

¹H NMR δ (CDCl₃) 4.62-4.70 (1/2H, m), 4.50-4.60 (1/2H, m), 3.38 -3.65 (2H, m), 3.15 (3H, 2s), 2.98 (3H, 2s), 1.90-2.21 (2H, m), 1.78-1.90 (2H, m), 1.40-1.42 (9H, 2s).

20

ii) N,N-Dimethyl-L-prolinamide hydrochloride

To the subtitle product of step i) (5.0g) was added hydrochloric acid (20ml, 4M). The mixture was stirred at room temperature for 3 h before the solvent was evaporated to afford the subtitle compound as a colourless solid. Yield: 3.50g

25 ¹H NMR δ (DMSO) 9.90 (1H, s, br), 8.40 (1H, s, br), 4.50-4.56 (1H, m), 3.00 -3.30 (2H, m), 3.04 (3H, s), 2.90 (3H, s), 2.21-2.44 (1H, m), 1.71-1.95 (3H, m).

iii) 1-(Aminosulfonyl)-N,N-dimethyl-L-prolinamide

To a solution of the subtitle product of step ii) (3.70g) in dioxane (50ml) was added
30 triethylamine (2.02g) and sulfamide (9.96g). The mixture was heated at reflux for 3 days before the reaction mixture was cooled, filtered and washed with methanol (50ml). The solvent was evaporated and the residue purified by column chromatography (2% methanol / EtOAc) to afford the subtitle compound as an oil. Yield: 1.70g.

MS APCI(+ve) 222[M+H]⁺

¹H NMR δ (DMSO) 6.63 (2H, bs), 4.58-4.61 (1H, m), 3.18 -3.34 (2H, m), 3.04 (3H, s), 2.80 (3H, s), 2.00-2.10 (1H, m), 1.68-1.92 (3H, m).

5 iv) 1-({[2-(Benzylthio)-6-chloropyrimidin-4-yl]amino}sulfonyl)-*N,N*-dimethyl-*L*-prolinamide


60% Sodium hydride (0.42g) was added to a solution of the subtitle product of step iii) (1.0g) in DMF (10ml) at 0°C. This mixture was stirred for 1h before the addition of the subtitle product of Example 19 step ii) (1.60g) in DMF (5ml) was added dropwise. The mixture was

10 allowed to reach room temperature and stirred for 3 days. To the mixture was added aqueous hydrochloric acid (1M) and extracted with EtOAc (2x100ml). The organic layer was washed with H₂O (2x50ml), the organic layer collected and concentrated causing the subtitle compound to precipitate out. This was filtered and washed with EtOAc (20ml). Yield: 1.10g. MS APCI(+ve) 455[M+H]⁺

15 ¹H NMR δ (CDCl₃) 7.42-7.46 (2H, m), 7.20-7.32 (3H, m), 7.02 (1H, s), 4.85-4.89 (1H, m), 4.32 (2H, s), 3.55- 3.60 (2H, m), 3.05 (3H, s), 3.02 (3H, s), 2.33-2.4 (1H, m), 1.94-2.13 (3H, m).

Example 26

20 1-{{[2-(Benzylthio)-6-{{(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl]amino}sulfonyl}-*N,N*-dimethyl-*D*-prolinamide



25 To the subtitle product of step iv) (1.0g) was added *R*-alaninol (3 ml) and the mixture heated at 80-90°C for 4 days. The reaction mixture was treated with EtOAc (100ml) and acidified with aqueous hydrochloric acid (1M, 100ml). The aqueous was separated and evaporated to give a residue which was purified by column chromatography (10% methanol / EtOAc) to

-56-

give the title compound as a white solid. This material was further purified by reverse phase HPLC (90% to 5% 0.02M ammonium hydroxide / acetonitrile). Yield: 70mg.

MS APCI(+ve) 495[M+H]⁺

¹H NMR δ (DMSO) 7.36-7.41 (2H, m), 7.20-7.30 (3H, m), 5.70 (1H, s), 5.23 (1H, bd), 4.70
5 (1H, t), 4.27-4.37 (2H, m), 3.15-3.31 (4H, m), 3.04 (3H, s), 2.80 (3H, s), 2.00-2.10 (1H, m), 1.80-2.00 (1H, m), 1.62-1.80 (2H, m), 1.07 (3H, d).

The intermediates for the above compound were prepared as follows:

i) 1-(Benzyloxycarbonyl)-*N,N*-dimethyl-*D*-prolinamide

10 The subtitle compound was prepared according to the method of Example 25 step i) using 1-(benzyloxycarbonyl)-*D*-proline (3.90g) in DCM (50ml). Yield: 4.45g.

MS APCI(+ve) 276[M+H]⁺

ii) *N,N*-Dimethyl-*D*-prolinamide

15 To the subtitle compound of step i) (4.45g) was added palladium hydroxide (0.20g) and methanol (50ml). The mixture was stirred under a hydrogen atmosphere (4 bar) at room temperature for 3h before the catalyst was filtered through celite and the solvent was evaporated to afford the subtitle compound as a solid. Yield: 2.25g.

¹H NMR δ (CDCl₃) 3.81-3.90 (1H, m), 3.13-3.22 (1H, m), 3.00 (3H, s), 2.98 (3H, s), 2.74-2.87
20 (1H, m), 2.50 (1H, bs), 2.05-2.20 (1H, m), 1.59-1.90 (3H, m).

iii) 1-(Aminosulfonyl)-*N,N*-dimethyl-*D*-prolinamide

To a solution of the subtitle compound of step ii) (2.20g) in dioxane (25ml) was added sulfamide (7.20g) and the mixture heated at reflux for 45h. The reaction mixture was cooled
25 and adsorbed on to silica gel and then purified by column chromatography (5% methanol / EtOAc) to afford the subtitle compound as an oil. Yield: 1.6g.

MS APCI(+ve) 222[M+H]⁺

¹H NMR δ (DMSO) 6.63 (2H, bs), 4.58-4.61 (1H, m), 3.18-3.40 (2H, m), 3.04 (3H, s), 2.80 (3H, s), 2.00-2.10 (1H, m), 1.70-2.00 (3H, m).

30

iv) 1-([2-(Benzylthio)-6-chloropyrimidin-4-yl]amino)sulfonyl)-*N,N*-dimethyl-*D*-prolinamide

-57-


60% Sodium hydride (0.42g) was added to a solution of the subtitle product of step iii) (1.50g) in DMF (20ml) at 0°C. This mixture was stirred for 1h and then a solution of the subtitle product of Example 19 step ii) (1.60g) in DMF (5ml) was added dropwise. The mixture was allowed to reach room temperature and stirred there for 2 days. To the mixture
5 was added aqueous hydrochloric acid (100ml) and the resulting precipitate was filtered. This solid was then stirred with EtOAc (200ml) and filtered to afford the subtitle compound as a solid. Yield: 2.0g.

MS APCI(+ve) 456[M+H]⁺

¹H NMR $\delta_{(\text{DMSO}-d_6)}$ 7.40-7.50 (2H, m), 7.20-7.40 (2H, m), 7.21-7.30 (1H, m) 6.70 (1H, s),
10 5.10-5.20 (1H, m), 4.32 (2H, s), 3.20-3.40 (2H, m), 3.04 (3H, s), 2.80 (3H, s), 2.00-2.10 (1H, m), 1.80-1.88 (1H, m), 1.70-1.88 (2H, m).

Example 27

N-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino)-
15 pyrimidin-4-yl)methanesulfonamide



Methanesulfonyl chloride (0.15ml) was added to a solution of the subtitle product of step iii) (0.40g) and *N,N*-diisopropylethylamine (0.53ml) in DCM (15ml) and stirring maintained for
20 30min. A further portion of methanesulfonyl chloride (0.15ml) and *N,N*-diisopropylethylamine (0.53ml) were added and stirring maintained for a further 30min. The reaction solution was extracted with H₂O (2 x 20ml) and the organics dried (MgSO₄) and concentrated to yield a brown oil. The residue was diluted in THF (7ml) and treated with aqueous sodium hydroxide solution (1.5ml of a 2.5M solution) for 1h at room temperature.
25 The reaction mixture was acidified with 2M hydrochloric acid solution to pH 1 and stirring maintained for 1h before the mixture was neutralised with sodium hydroxide solution, concentrated onto silica gel and purified by column chromatography (Et₂O then EtOAc). The crude product was then purified by reverse phase HPLC (acetonitrile / 0.02M ammonium

-58-

hydroxide (90% to 5% aqueous phase)) to yield the title compound as a white solid. Yield: 0.12g.

MS APCI(+ve) 421 [M+H]⁺

¹H NMR δ (DMSO) 7.57 (1H, t), 7.48 (1H, t), 7.26 (1H, s), 7.17 (1H, t), 5.85 (1H, s), 4.71 (1H, s), 4.38 (2H, s), 3.98 (1H, s), 3.43 - 3.24 (2H, m), 3.20 (3H, s), 1.05 (3H, d).

The intermediates for the above compound were prepared as follows:

i) 6-Amino-2-[(3-chloro-2-fluorobenzyl)thio]pyrimidin-4(3H)-one

The subtitle compound was prepared according to the procedure of Example 1 step i) treating 4-amino-6-hydroxy-2-mercaptopyrimidine monohydrate (8.4g) with 3-chloro-2-fluorobenzyl bromide (11.0g) to afford the subtitle compound as a white solid. Yield: 14.1g.

MS APCI(+ve) 286 [M+H]⁺

ii) 6-Chloro-2-[(3-chloro-2-fluorobenzyl)thio]pyrimidin-4-amine

The subtitle compound was prepared from the product of step i) (2.00g) according to the procedure of Example 1 step ii) to afford the subtitle product as a green foam which was used directly in the next step.

MS: APCI(+ve) 304 [M+H]⁺


iii) N-((1R)-2-[[tert-Butyl(dimethyl)silyl]oxy]-1-methylethyl)-2-[(3-chloro-2-fluorobenzyl)thio]pyrimidine-4,6-diamine

N,N-Diisopropylethylamine (5.2ml) was added to a solution of (R)-alaninol (2.56ml) and the subtitle product of step ii) in NMP (35ml) and stirred at 100°C for 24h and then 140°C for 24h. After cooling to ambient temperature imidazole (2.60g) and a solution of tert-butyl dimethylsilyl chloride (2.60g) in THF (10ml) were added and stirring maintained for 1h. The volatiles were removed *in vacuo* and the residue was purified by column chromatography (1:1 Et₂O / iso-hexane) to afford the subtitle compound as a yellow oil. Yield: 1.70g.

MS: APCI(+ve) 457 [M+H]⁺

Example 28

N-(2-[(2,3-Dichlorobenzyl)thio]-6-[(1R)-2-hydroxy-1-methylethyl]amino)pyrimidin-4-yl)methanesulfonamide



The title compound was synthesised from the subtitle product of step iii) (0.4g) according to the procedure of Example 27 to afford the title compound as a white solid.

5 Yield: 0.15g.

MS APCI(+ve) 437 $[M+H]^+$

^1H NMR δ (DMSO) 10.55 (1H, s), 7.66 (1H, d), 7.55 (1H, dd), 7.31 (1H, t), 7.26 (1H, s), 5.78 (1H, s), 4.71 (1H, s), 4.46 (2H, s), 3.99 (1H, s), 3.41 - 3.24 (2H, m), 3.20 (3H, s), 1.05 (3H, d).

10 The intermediates for the above compound were prepared as follows:

i) 6-Amino-2-[(2,3-dichlorobenzyl)thio]pyrimidin-4(3H)-one

The subtitle compound was prepared according to the procedure of Example 1 step i) treating 4-amino-6-hydroxy-2-mercaptopyrimidine monohydrate (2.22g) with 2,3-dichlorobenzyl bromide (3.30g) to afford the subtitle compound as a white solid. Yield: 3.25g.

15 MS APCI(+ve) 302 $[M+H]^+$

ii) 6-Chloro-2-[(2,3-dichlorobenzyl)thio]pyrimidin-4-amine

The subtitle compound was prepared from the product of step i) (2.00g) according to the procedure of Example 1 step ii) to afford the subtitle product as a green foam which was used

20 directly in the next step.

MS: APCI(+ve) 320 $[M+H]^+$


iii) N-((1R)-2-[(tert-Butyl(dimethyl)silyl)oxy]-1-methylethyl)-2-[(2,3-dichlorobenzyl)thio]pyrimidine-4,6-diamine

25 The subtitle compound was synthesised from the subtitle product of step ii) according to the procedure of Example 27 step iii). Yield: 0.4g.

MS: APCI(+ve) 473 $[M+H]^+$

Example 29

***N*-(2-[(3-Chlorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)methanesulfonamide**



5

The title compound was prepared from the subtitle product of step iii) (0.4g) according to the procedure of Example 27 as a white solid. Yield: 71mg.

MS APCI(+ve) 403 $[M+H]^+$

^1H NMR δ (DMSO) 7.52 - 7.17 (4H, m), 5.78 (1H, s), 4.70 (1H, t), 4.32 (2H, dd), 3.97 (1H, s),
10 3.44 - 3.24 (2H, m), 3.18 (3H, s), 1.06 (3H, d).

The intermediates for the above compound were prepared as follows:

i) 6-Amino-2-[(3-chlorobenzyl)thio]pyrimidin-4(3*H*)-one

The subtitle compound was prepared according to the procedure of Example 1 step i) treating 4-amino-6-hydroxy-2-mercaptopyrimidine monohydrate (10.00g) with 3-chlorobenzyl

15 chloride (9.98g) to afford the subtitle compound as a white solid. Yield: 15.00g.

MS APCI(+ve) 268 $[M+H]^+$

ii) 6-Chloro-2-[(3-chlorobenzyl)thio]pyrimidin-4-amine

The subtitle compound was prepared from the product of step i) (2.00g) according to the
20 procedure of Example 1 step ii) to afford the subtitle product as a green foam which was used directly in the next step.

MS: APCI(+ve) 286 $[M+H]^+$


25 iii) *N*-((*1R*)-2-[[*tert*-Butyl(dimethyl)silyl]oxy]-1-methylethyl)-2-[(3-chlorobenzyl)thio]-pyrimidine-4,6-diamine

The subtitle compound was synthesised from the subtitle product of Example 29 step ii) according to the procedure of Example 27 step iii). Yield: 0.4g.

MS: APCI(+ve) 439 $[M+H]^+$

Example 30

***N*-(2-[(2-Fluoro-4-methoxybenzyl)thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino)-pyrimidin-4-yl)methanesulfonamide**



5

The title compound was synthesised from the subtitle product of Example 30 step iii) (0.4g) according to the procedure of Example 27 as a white solid. Yield: 35mg.

MS APCI(+ve) 403 [M+H]⁺

¹H NMR δ_(DMSO) 7.45 (1H, t), 7.21 (1H, s), 6.81 (1H, dd), 6.73 (1H, dd), 5.77 (1H, s) 4.71
10 (1H, t), 4.27 (2H, s), 3.74 (3H, s), 3.44 - 3.21 (2H, m), 3.20 (3H, s), 1.08 (3H, d).

The intermediates for the above compound were prepared as follows:

i) 6-Amino-2-[(2-fluoro-4-methoxybenzyl)thio]pyrimidin-4(3*H*)-one

The subtitle compound was prepared according to the procedure of Example 1 step i) treating
15 4-amino-6-hydroxy-2-mercaptopyrimidine monohydrate (1.86g) with 2-fluoro-4-methoxybenzyl chloride (2.00g) to afford the subtitle compound as a white solid. Yield: 2.64g.

MS APCI(+ve) 282 [M+H]⁺

ii) 6-Chloro-2-[(2-fluoro-4-methoxybenzyl)thio]pyrimidin-4-amine

20 The subtitle compound was prepared from the product of Example 30 step i) according to the procedure of Example 1 step ii) to afford the subtitle product as a green foam which was used directly in the subsequent step.

MS: APCI(+ve) 300 [M+H]⁺


25 iii) *N*-((1*R*)-2-[(*tert*-Butyl-(dimethyl)silyl)oxy]-1-methylethyl)-2-[(2-fluoro-4-methoxybenzyl)thio]pyrimidine-4,6-diamine

The subtitle compound was synthesised from the subtitle product of step ii) according to the procedure of Example 27 step iii). Yield: 0.4g.

MS: APCI(+ve) 453 [M+H]⁺

Example 31

***N*-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)piperazine-1-sulfonamide**



The subtitle product of step ii) (0.7g) was diluted in (*R*)-alaninol (4ml) and heated at 80°C for 48h. The product was purified through a plug of silica gel (1:1 EtOAc / isohexane then 90:9:1 EtOAc / methanol / *N,N*-diethylisopropylamine). The crude product was diluted in 1,4-dioxane (40ml) and treated with HCl/1,4-dioxane (0.5ml of a 4M solution) for 1h before concentrating *in vacuo* and purifying the crude product by reverse phase HPLC (acetonitrile / 0.02M ammonium hydroxide (90% to 5% aqueous phase)) to yield the title product as a white solid. Yield: 49mg.

MS: APCI(+ve) 491 [M+H]⁺

15 ¹H NMR δ_(DMSO) 7.59 (1H, t), 7.46 (1H, t), 7.15 (1H, t), 7.06 (1H, s), 5.85 (1H, s), 4.69 (1H, t), 4.36 (2H, t), 3.89 (1H, s), 3.42 - 3.23 (2H, m), 3.05 (4H, m), 2.71 (4H, m), 1.05 (3H, d).

The intermediates for the above compound were prepared as follows:

i) *tert*-Butyl 4-(aminosulfonyl)piperazine-1-carboxylate

20 A mixture of *tert*-butyl-4-piperazine-1-carboxylate (1.47g) and sulfamide (5.25g) in 1,4-dioxan (50ml) were heated at reflux for 48h. The volatiles were removed *in vacuo* and the residue partitioned between EtOAc (100ml) and H₂O (100ml). The organics were recovered and the aqueous extracted further with EtOAc (3 x 100ml). The organics were then combined, dried (MgSO₄) and concentrated. The crude material was triturated with diethyl ether and filtered to provide the subtitle compound as a white solid. Yield: 1.91g

¹H NMR δ_(DMSO) 6.80 (2H, s), 3.40 (4H, t), 2.90 (4H, t), 1.41 (9H, s).

ii) 2-[(3-Chloro-2-fluorobenzyl)thio]pyrimidine-4,6-diol

-63-

The subtitle compound was prepared by the method of Example 19 step i) using 2-mercaptopyrimidine-4,6-diol (20.0g) and 3-chloro-2-fluorobenzyl bromide to afford the subtitle compound as a white solid. Yield: 36.2g.

MS: APCI(+ve) 287 [M+H]⁺

5

iii) 4,6-Dichloro-2-[(3-chloro-2-fluorobenzyl)thio]pyrimidine

N,N-Dimethylaniline (50ml) was added to a slurry of the the subtitle product of step ii) (36.2g) in phosphorus oxychloride (200ml) and heated at reflux for 10h. The reaction was allowed to cool and the excess phosphorus oxychloride removed *in vacuo* before pouring onto
10 ice. This mixture was extracted with EtOAc (400ml) and washed with brine (2x100ml), the organics collected, dried (MgSO₄) and concentrated *in vacuo* to afford the crude product. This crude product was purified by flash chromatography (EtOAc /isohexane (2 to 5%)) to yield the subtitle compound as an oil. Yield: 6.2g.

MS: APCI(-ve) 322 [M-H]⁺

15 iv) *N*-{6-Chloro-2-[(3-chloro-2-fluorobenzyl)thio]pyrimidin-4-yl}piperazine-1-sulfonamide


60% Sodium hydride (0.22g) was added to a solution of the subtitle product of step i) (1.78g) in DMF (10ml) at 0°C. The reaction was removed from the cooling bath and stirred for 1h before adding the subtitle product of step iii) (1.46g) as a solution in DMF (5ml). The
20 reaction was stirred overnight before concentrating *in vacuo*. The residue was partitioned between H₂O (50ml) and EtOAc (50ml) and the organics recovered, dried (MgSO₄) and concentrated *in vacuo*. The crude product was purified by flash column chromatography (20 % then 30 % EtOAc / iso-hexane) to yield the subtitle compound as a white solid. Yield: 0.7g.
¹H NMR δ_(DMSO) 11.64 (1H, s), 7.63 - 7.47 (2H, m), 7.19 (1H, t), 6.70 (1H, s), 4.47 (2H, s),
25 3.35 (8H, s), 1.40 (9H, s).

Example 32

***N*-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[(1*R*)-1-(hydroxymethyl)propyl]amino}-pyrimidin-4-yl)piperazine-1-sulfonamide**

30

-64-



The subtitle product of step iv) (0.27g) was diluted in trifluoroacetic acid (2ml) and stirred for 20min before removing the volatiles *in vacuo*. The residue was diluted in DCM (20ml) and the volatiles again removed *in vacuo*. The title product was purified by reverse phase HPLC (acetonitrile / 0.02M ammonium hydroxide (90% to 5% aqueous phase)) to yield the title compound as a white solid. Yield: 0.11g.

MS: APCI(+ve) 505 $[M+H]^+$

^1H NMR $\delta_{(\text{DMSO})}$ 7.59 (1H, t), 7.47 (1H, t), 7.15 (1H, t), 7.10 (1H, s), 5.89 (1H, s), 4.64 (1H, s), 4.36 (2H, s), 3.43 - 3.24 (2H, m), 3.08 (4H, m), 2.72 (4H, d), 1.60 (1H, m), 1.36 (1H, m), 0.82 (3H, t).

The intermediates for the above compound were prepared as follows:

i) ***tert*-Butyl-4-[(6-chloro-2-[(3-chloro-2-fluorobenzyl)thio]pyrimidin-4-yl){[2-(trimethylsilyl)ethoxy]methyl}amino)sulfonyl]piperazine-1-carboxylate**

60% Sodium hydride (44mg) was added to a solution of the subtitle product of Example 31 step iii) (0.40g) in DMF (6ml) at 0°C. Stirring was maintained for 10min before the addition of [2-(chloromethoxy)ethyl](trimethyl)silane. The reaction was removed from the cooling bath and stirred for 2h before the reaction was partitioned between H₂O (30ml) and EtOAc (50ml). The aqueous was extracted further with EtOAc (2 x 50ml), the organics combined, dried (MgSO₄) and concentrated to afford the subtitle product as a colourless oil. Yield: 0.60g.

MS: APCI(+ve) 682 $[M+H]^+$

ii) ***tert*-butyl-4-[(2-[(3-chloro-2-fluorobenzyl)thio]-6-[(1*R*)-1-(hydroxymethyl)propyl]amino)pyrimidin-4-yl){[2-(trimethylsilyl)ethoxy]methyl}amino)sulfonyl]piperazine-1-carboxylate**

-65-


The subtitle product of step i) (0.60g) was diluted in (2*R*)-2-aminobutan-1-ol (2ml) and heated at 80°C for 18h. The subtitle compound was recovered as a colourless oil by flash column chromatography (EtOAc / iso-hexane mixtures). Yield: 0.40g.

MS: APCI(+ve) 735 [M+H]⁺

5

Example 33

N'-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)-*N,N*-dimethylsulfamide



10

The subtitle product of step i) (0.7g) was diluted in (*R*)-alaninol (4ml) and heated at 80°C for 48h before partitioning between EtOAc / 1M hydrochloric acid. The aqueous was further extracted with EtOAc (3x), the organics combined, washed with brine, dried (MgSO₄) and

15 concentrated. The product was purified by column chromatography (25%, 40% EtOAc / iso-hexane then EtOAc) to yield the title compound as a colourless oil. Yield: 0.13g.

MS: APCI(+ve) 450 [M+H]⁺

¹H NMR δ(DMSO) 10.43 (1H, s), 7.57 (1H, t), 7.47 (1H, dd), 7.29 (1H, s), 7.16 (1H, t), 5.87 (1H, s), 4.70 (1H, s), 4.37 (2H, t), 4.04 (1H, s), 3.43 - 3.24 (2H, m), 2.77 (6H, s), 1.05 (3H, d).

20

The intermediates for the above compound were prepared as follows:

i) *N'*-{6-chloro-2-[(3-chloro-2-fluorobenzyl)thio]pyrimidin-4-yl}-*N,N*-dimethyl-sulfamide

60% Sodium hydride (0.48g) was added to a solution of *N,N*-dimethylsulfamide (1.05g) in DMF (15ml) and stirring maintained for 10min before the addition of the subtitle product of

25 Example 31 step iii) (3.56g). The reaction was stirred for 18h before the reaction was partitioned between H₂O (30ml) and EtOAc (50ml). The aqueous was extracted further with EtOAc (2 x 50ml) the organics combined, washed with H₂O (100ml), brine (50ml), dried (MgSO₄) and concentrated. The crude product was purified by flash column chromatography

-66-


(20 % then 30 % EtOAc / isohexane) to yield the subtitle compound as a white solid. Yield: 1.57g.

^1H NMR $\delta_{(\text{DMSO})}$ 11.49 (1H, s), 7.56 (1H, t), 7.51 (1H, t), 7.19 (1H, t), 6.69 (1H, s), 4.44 (2H, s), 2.83 (6H, s).

5

Example 34

***N*-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-2-(dimethylamino)ethanesulfonamide**



10

The subtitle product of step iv) (0.7g) was diluted in (*R*)-alaninol (2.25ml) and heated at 80°C for 48h. The product was purified by reverse phase HPLC (acetonitrile / 0.02M ammonium hydroxide (90% to 5% aqueous phase)) to yield the title compound as a white solid. Yield:

15 46mg.

MS: APCI(+ve) 478 $[\text{M}+\text{H}]^+$

^1H NMR $\delta_{(\text{DMSO})}$ 7.6 (1H, t), 7.43 (1H, t), 7.15 (1H, t), 6.64 (1H, br. s), 5.54 (1H, s), 4.35 (2H, s), 3.80 (1H, s), 3.41-3.11 (6H, m), 2.09 (6H, s) and 1.08 (3H, d).

20 The intermediates for the above compound were prepared as follows:

i) Sodium 2-(dimethylamino)ethanesulfonate

Sodium ethylenesulfonate (18.2ml, 25% aqueous solution w/w) and dimethylamine (4.43ml, 40% aqueous solution w/w) were heated in a sealed finger tube at 105°C for 48h before concentrating *in vacuo* to yield the subtitle compound as a white solid. Yield: 9.80g.

25 ^1H NMR $\delta_{(\text{CD}_3\text{OD})}$ 3.05 (2H, m), 2.84 (2H, m) and 2.30 (6H, s).

ii) 2-(Dimethylamino)ethanesulfonyl chloride

-67-

Phosgene (12ml, 1M in toluene) was added to a solution of the subtitle product of step i) in DCM (20ml) and DMF (2.5ml) and stirred for 3h before concentrating *in vacuo* to afford the subtitle compound as an oil that was used directly in the next step.

^1H NMR $\delta_{(\text{CD}_3\text{OD})}$ 3.30 (2H, m), 2.80 (2H, m) and 2.31 (6H, s).

5

iii) 2-(Dimethylamino)ethanesulfonamide

Ammonia was cautiously added to a cooled solution of the subtitle product of step ii) in THF (50ml) and stirring maintained until all of the ammonia had evaporated before filtering off the product as a white solid. Yield: 0.88g.

10 ^1H NMR $\delta_{(\text{CDCl}_3)}$ 3.20 (2H, t), 2.86 (2H, m) and 2.30 (6H, s).


iv) *N*-{6-Chloro-2-[(3-chloro-2-fluorobenzyl)thio]pyrimidin-4-yl}-2-(dimethylamino)ethanesulfonamide

The subtitle compound was prepared by the method of Example 33 step i) from the reaction
15 of the subtitle product of step iii) (0.8g) and the subtitle product of Example 31 step iii) (1.1g) and used directly in the following step. Yield 0.7g

^1H NMR $\delta_{(\text{CDCl}_3)}$ 7.48 (1H, t), 7.31 (1H, t), 7.02 (1H, t), 6.87 (1H, s), 4.38 (2H, s), 3.21 (2H, m), 2.83 (2H, m), 2.32 (3H, s), 2.25 (3H, s).

20 **Example 35**

***N*-(2-[(3-chloro-2-fluorobenzyl)thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino)pyrimidin-4-yl)-4-methylpiperazine-1-sulfonamide**



25 A solution of the product from step iv) (1.5g) in (*R*)-alaninol (3ml) was heated at 80°C for 4 days. The resulting mixture was diluted with acetonitrile (5ml) and purified by reverse phase HPLC (50% to 5% 0.02M ammonium hydroxide / methanol) to yield the title compound as a white solid. Yield: 1.0g.

MS APCI(+ve) 506 $[\text{M}+\text{H}]^+$

-68-

^1H NMR $\delta_{(\text{DMSO})}$ 10.60 (1H, bd), 7.60 (1H, m), 7.55 (1H, m), 7.47 (1H, bs), 7.19 (1H, t), 5.88 (1H, s), 4.72 (1H, t), 4.37 (2H, s), 3.40 (1H, m), 3.30 (2H, m), 3.10 (4H, m), 2.30 (4H, m), 2.10 (3H, s), 1.05 (3H, d).

5 The intermediates for this compound were prepared as follows:

i) 4-Methyl-1-piperazinesulfonamide

N-Methyl piperazine (5ml) and sulfamide (11.26g) in 1,4-dioxane (100ml) were heated at reflux for 48h. The solvent was evaporated under reduced pressure and the resulting solid dissolved in a mixture of methanol and water and applied to a plug of SCX-silica followed by
10 further elution with aqueous methanol. This was then followed by elution with 1M ammonia in methanol (x4) and these fractions collected and solvent evaporated to dryness leaving a white solid. This was triturated with Et_2O and filtered leaving the subtitle compound as a white solid. Yield: 5.0g.

^1H NMR $\delta_{(\text{DMSO})}$ 6.75 (2H, s), 2.90 (4H, m), 2.40 (4H, m), 2.20 (3H, s).

15

ii) 2-[(3-Chloro-2-fluorobenzyl)thio]-4,6-pyrimidinediol

To a slurry of 2-mercapto-4,6-pyrimidinediol (64.6g) in ethanol (387ml) and H_2O (387ml) was added sodium hydroxide (18g) in H_2O (82ml) causing almost a complete solution to form. 2-Fluoro-3-chloro-benzylbromide (100g) in ethanol (82ml) was then added dropwise
20 over 30min causing a thick precipitate to form. Stirring was continued for a further 4h and the whole then cooled to 0°C before filtering the white solid formed and then washing with water. The collected solid was dried *in vacuo* at 50°C for 48h leaving the subtitle compound as a white solid. Yield: 125.7g.

MS APCI(+ve) 287 $[\text{M}+\text{H}]^+$

25

iii) 4,6-Dichloro-2-[(3-chloro-2-fluorobenzyl)thio]-pyrimidine

The subtitle product from step ii) (125.67g) was added slowly to phosphorus oxychloride (1L) over 10min with stirring. After the addition was complete, *N,N*-dimethylaniline (92ml) was added portionwise with care over 10min causing a complete solution to form. This was then
30 stirred at 120°C for 15h. The cooled reaction mixture was then poured onto crushed ice and extracted with EtOAc (x3). The organic phases were combined, dried (MgSO_4) and the solvent evaporated leaving a brown oil which was purified by column chromatography (2% EtOAc / iso-hexane) to afford the subtitle compound as a white solid. Yield: 113g.

-69-

^1H NMR $\delta_{(\text{CDCl}_3)}$ 7.41(1H, m), 7.30(1H, m), 7.0 (2H, m + s), 4.40(2H, s).

iv) ***N*-[6-Chloro-2-[(3-chloro-2-fluorobenzyl)thio]-4-pyrimidinyl]-4-methyl-1-piperazinesulfonamide**


- 5 To a solution of the subtitle product of step i) (2.0g) in dry DMF (20ml) at 0°C was added 60% sodium hydride (0.9g) portionwise over 5min. After further stirring at 0°C for 30min the subtitle product from step iii) (3.6g) in DMF (10ml) was added and the whole allowed to stir at room temperature for 24h. The reaction mixture was carefully quenched with aqueous 2M hydrochloric acid until pH 7.4. The solvent was evaporated to dryness and the residue
- 10 dissolved in a mixture of methanol / EtOAc and applied to an SCX column followed by further elution with EtOAc. This was then followed by elution with EtOAc / triethylamine mixtures and these fractions evaporated to dryness. The subtitle compound was obtained pure by trituration with Et₂O and filtration, leaving a white solid. Yield: 1.5g.

MS APCI(+ve) 467 [M+H]⁺

15

Example 36

***N*-[2-[(3-Chloro-2-fluorobenzyl)thio]-6-[(2-hydroxy-1-methylethyl)amino]-4-pyrimidinyl]-4-morpholinesulfonamide**



The title compound was prepared from the product of step ii) (1.0g) and (*R*)-alaninol (5ml) according to the procedure used in Example 35 as a white solid. Yield: 0.68g.

MS APCI(+ve) 493 [M+H]⁺

- 25 ^1H NMR $\delta_{(\text{DMSO})}$ 10.60 (1H, bs), 7.60 (1H, m), 7.50 (1H, m), 7.30 (1H, bs), 7.19 (1H, m), 5.90 (1H, s), 4.70 (1H, m), 4.20 (2H, s), 4.0 (1H, m), 3.80 (4H, m), 3.40 (2H, m), 3.10 (4H, m), 1.10 (3H, d).

The intermediates for this compound were prepared as follows

-70-

i) 4-Morpholinesulfonamide

Morpholine (5ml) and sulfamide (11g) in 1,4-dioxane (100ml) were heated at reflux for 48h. The solvent was evaporated under reduced pressure and the resulting solid partitioned between EtOAc and water. The organic phase was collected and the aqueous phase was further extracted with EtOAc (x4). The combined organic extracts were dried (MgSO₄) and the solvent removed *in vacuo*. The solid residue was triturated with Et₂O and filtered to give the subtitle compound as a white crystalline solid. Yield: 2.1g.

¹H NMR $\delta_{(DMSO)}$ 6.82 (2H, s), 3.66 (4H, m), 2.90 (4H, m).


10 **ii) N-[6-chloro-2-[(3-chloro-2-fluorobenzyl)thio]-4-pyrimidinyl]-4-morpholinesulfonamide**

To a solution of the product of step i) (2.1g) in dry DMF (20ml) at 0°C under nitrogen was added 60% sodium hydride (1.1g) portionwise over 5min. After further stirring at 0°C for 30min the product from Example 31 step iii) (4g) in DMF (10ml) was added and the whole allowed to further stir at room temperature for 24h. The reaction mixture was carefully quenched with aqueous 2M hydrochloric acid until pH 7.4. The solvent was evaporated to dryness and the residue partitioned between EtOAc and brine. The organic phase was collected, dried (MgSO₄) and the solvent removed *in vacuo* to afford the subtitle compound as a gummy solid. Yield: 1.3g.

20 MS APCI(+ve) 454 [M+H]⁺

Example 37

N-[2-[[[(3-Chloro-2-fluorophenyl)methyl]thio]-6-[(2-hydroxy-1-methylethyl)amino]-4-pyrimidinyl]-1,2-dimethyl-1H-imidazole-4-sulfonamide



25

To a solution of the product from Example 27 step iii) (1.3g) in dry pyridine (10ml) and 4-*N,N*-dimethylaminopyridine (0.48g) under nitrogen was added 1,2-dimethyl-1H-imidazole-4-

-71-


sulfonyl chloride (1.0g). The reaction mixture was heated at 55°C for 5 days. The cooled reaction mixture was partitioned between EtOAc and 1M hydrochloric acid. The organic phases were collected, dried (MgSO₄) and the solvent evaporated to dryness. The resulting gum was dissolved in acetonitrile (15ml) and treated with 2M hydrochloric acid (5ml) and the whole stirred at room temperature for 15 min. The reaction mixture was evaporated to dryness leaving a gummy residue. Purification by column chromatography (DCM / methanol / acetic acid 190:10:1) afforded the title compound as a white solid. Yield: 1.0g.

MS APCI(+ve) 501 [M+H]⁺

¹H NMR δ_(DMSO) 7.85 (1H, bs), 7.50 (2H, m), 7.18 (1H, m), 5.91(1H, m), 4.36 (2H, s), 3.60 (3H, s), 3.30 (2H, m), 2.30 (3H, s), 1.10 (3H, d).

Example 38

N-(2-[(2,3-Difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)piperazine-1-sulfonamide



The subtitle product of step i) (2.0g) was diluted in DCM / trifluoroacetic acid (10:1) for 1h before concentrating *in vacuo* and purifying the crude product by reverse phase HPLC (acetonitrile / 0.02M ammonium hydroxide (90% to 5% aqueous phase)) to yield the title compound as a white solid. Yield: 6mg.

MS: APCI(+ve) 475 [M+H]⁺

¹H NMR δ_(DMSO) 7.41 (1H, m), 7.31 (1H, m), 7.14 (1H, m), 5.87 (1H, s), 4.70 (1H, t), 4.38 (2H, s), 3.93 (1H, br. s), 3.44-3.26 (4H, m), 2.81-2.67 (5H, m), 2.42 (1H, m), 1.05 (3H, d).

The intermediates for the above compound were prepared as follows:

i) **Piperazine-1-sulfonamide-4,6-dichloro-2-[(3-chloro-2-fluorobenzyl)thio]-pyrimidine**
60% Sodium hydride (0.26g) was added to a solution of the subtitle product of Example 31 step i) (3.11g) in DMF (10ml) at 0°C. The reaction was removed from the cooling bath and

-72-

stirred for 1h before adding the subtitle product of Example 39 step ii) (5.0g) as a solution in DMF (5ml). The reaction was stirred overnight before concentrating *in vacuo*. The residue was partitioned between H₂O (50ml) and EtOAc (50ml) and the organics recovered, dried (MgSO₄) and concentrated *in vacuo*. The crude product was purified by flash column chromatography (20 %, 25% then 30 % EtOAc / isohexane) to yield the intermediate as a white solid. This material was diluted in (*R*)-alaninol (12ml) and heated at 80°C for 72h. The residue was partitioned between H₂O (50ml) and EtOAc (50ml) and the organics recovered, dried (MgSO₄) and concentrated *in vacuo* to afford the subtitle compound as a white solid.


Yield: 2.0g.

¹H NMR δ_(DMSO) 7.15-6.93 (3H, m), 6.20 (1H, s), 5.00 (1H, d), 4.29 (2H, s), 3.71 (1H, dd), 3.57 (1H, dd), 3.42 (4H, t), 3.20 (4H, t), 1.46 (9H, s), 1.21 (3H, d).

Example 39

N-(2-[(2,3-Difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-

yl)azetidine-1-sulfonamide



A solution of the subtitle product of step iii) (0.5g) in (*R*)-alaninol (1.2ml) was heated at 80°C for 18h before partitioning between EtOAc and H₂O. The organics were recovered, dried (MgSO₄) and concentrated. The residue was purified by column chromatography (1:90:109 AcOH / EtOAc / iso-hexane) before treating the crude material with trifluoroacetic acid (2ml) and stirring for 12min then quenching the reaction by the addition of 1M sodium hydroxide solution to pH >10. Saturated ammonium chloride solution was then added to pH 4 and the organics recovered by extracting with EtOAc (3 x 20ml). The organics were dried (MgSO₄) and concentrated *in vacuo*. The crude material was purified by reverse phase HPLC (95% to 20% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 90mg.

MS APCI(+ve) 446 [M+H]⁺

-73-

^1H NMR $\delta_{(\text{CDCl}_3)}$ 7.24 - 7.21 (1H, m), 7.08 - 6.97 (2H, m), 6.01 (1H, s), 5.06 - 4.95 (1H, m), 4.35 (2H, s), 4.20 - 4.05 (1H, m), 3.98 (4H, t), 3.74 - 3.70 (1H, m), 3.60 - 3.56 (1H, m), 2.23 (2H, quin.), 1.22 (3H, d).

5 The intermediates for this compound were prepared as follows:

i) 2-[(2,3-Difluorobenzyl)thio]pyrimidine-4,6-diol

A solution of potassium hydroxide (5.67g) was added dropwise to a suspension of 2-mercaptopyrimidine-4,6-diol (14.56g) in DMF (78ml) and H_2O (39ml) and the mixture stirred for 30min. A solution of 2,3-difluorobenzyl bromide (20.86g) in THF (16ml) was then added
10 dropwise and the mixture stirred for 18h. The reaction was then cooled to 0°C and the precipitate was filtered and washed with H_2O (4 x 100ml) before drying *in vacuo* to afford the subtitle compound as a cream solid. Yield: 22.4g.

^1H NMR $\delta_{(\text{DMSO})}$ 7.74 (1H, s), 7.39 - 7.32 (2H, m), 7.21 - 7.15 (1H, m), 4.48 (2H, s).

15 ii) 4,6-Dichloro-2-[(2,3-difluorobenzyl)thio]pyrimidine

N,N-Dimethylaniline (10.3ml) was added to a slurry of the the subtitle product of step i) (10.0g) in phosphorus oxychloride (55ml) and the solution heated at reflux for 10h. The reaction was allowed to cool and excess phosphorus oxychloride was removed *in vacuo* before partitioning between Et_2O (110ml) and H_2O (275ml) and stirring for 1h. The layers
20 were separated and the organics concentrated *in vacuo* to afford the crude product. This crude product was purified by column chromatography (4% EtOAc / iso-hexane) to yield the subtitle compound as a white solid. Yield: 9.10g.

^1H NMR $\delta_{(\text{DMSO})}$ 7.74 (1H, s), 7.39-7.32 (2H, m), 7.21-7.15 (1H, m) 4.48 (2H, s).

25 iii) *N*-{6-Chloro-2-[(2,3-difluorobenzyl)thio]pyrimidin-4-yl}-*N*-{[2-(trimethylsilyl)ethoxy)methyl]azetidine-1-sulfonamide}

To a solution of the product of Example 40 step i) (0.33g) in dry DMF (4ml) at 0°C under nitrogen was added 60% sodium hydride (0.20g). The reaction was allowed to warm outside the cooling bath for 15min before recooling to 0°C and addition of the product from step ii)
30 (0.75g) in DMF (2ml) and the whole allowed to further stir at room temperature for 3h. The reaction was quenched with 2-(trimethylsilyl)ethoxymethyl chloride (0.86ml) and allowed to stir for 18h before removal of the volatiles *in vacuo* and partitioning of the residue between EtOAc (100ml) and H_2O (200ml). The aqueous was washed further with EtOAc (2 x 100ml)

-74-


and the organics combined, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography (1:22:177 AcOH / EtOAc / iso-hexane) to afford the subtitle compound as a colourless oil. Yield: 0.65g.

¹H NMR δ_(CDCl₃) 7.34 – 7.30 (1H, m), 7.19 (1H, s), 7.13 – 7.02 (2H, m), 5.42 (2H, s), 4.45 (2H, s), 4.06 (4H, t), 3.65 (2H, t), 2.27 (2h, quin.), 0.93 (2H, t), 0.00 (9H, s).

Example 40

N-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino)-pyrimidin-4-yl)azetidine-1-sulfonamide

10



A solution of the subtitle product of step vi) (0.5g) in trifluoroacetic acid (5ml) was stirred for 15min before the addition of 2M sodium hydroxide solution until pH >10. The aqueous was then extracted with Et₂O (20ml) before acidifying the aqueous to pH 4 with 2M hydrochloric acid and extracting with EtOAc (2 x 20ml). The EtOAc extracts were combined, dried (MgSO₄) and concentrated. The crude material was purified by reverse phase HPLC (90% to 5% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 10mg.

MS APCI(+ve) 462 [M+H]⁺

¹H NMR δ_(DMSO) 7.59 (1H, t), 7.46 (1H, t), 7.15 (1H, t), 5.90 (1H, s), 4.69 (1H, t), 4.37 (2H, s), 3.95 (1H, br.s), 3.81 (4H, m), 3.44 – 3.21 (2H, m), 2.12 (2H, m), 1.05 (3H, d).

The intermediates for this compound were prepared as follows:

25 i) Azetidine-1-sulfonamide

Azetidine (4.23g) was added to a solution of sulfamide (7.48g) in 1,4-dioxan (120ml) and heated at reflux for 24h. The volatiles were removed under reduced pressure and the residue suspended in refluxing CHCl₃ (500ml) for 10min before decanting. The residue was again

-75-

suspended in hot CHCl_3 (500ml) for 10min before decanting. The filtrates were combined and concentrated *in vacuo* to afford the subtitle product as a white solid. Yield: 4.1g.

^1H NMR $\delta_{(\text{DMSO})}$ 6.91 (2H, s), 3.74 (4H, t), 2.15 (2H, quin.).

5 **ii) *N*-[2-(Benzylthio)-6-chloropyrimidin-4-yl]-*N*-{[2-(trimethylsilyl)ethoxy]methyl}-azetidine-1-sulfonamide**

The subtitle compound was prepared as a yellow oil by the method of Example 32 step i) by reacting the subtitle product of step i) (4.6g) with the subtitle product of Example 19 step ii) (12.9). Yield: 12.9g.

10 MS APCI(+ve) 537 $[\text{M}+\text{H}]^+$

iii) *N*-(2-(Benzylthio)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-*N*-{[2-(trimethylsilyl)ethoxy]methyl}azetidine-1-sulfonamide

The subtitle compound was prepared as a yellow oil by the method of Example 32 step ii) by reacting the subtitle product of step ii) (12.2g) with (*R*)-alaninol (18ml). Yield: 11.3g.

MS APCI(+ve) 540 $[\text{M}+\text{H}]^+$

iv) *N*-(2-(Benzylsulfonyl)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-*N*-{[2-(trimethylsilyl)ethoxy]methyl}azetidine-1-sulfonamide

20 *m*-Chloroperbenzoic acid was added as a single portion to a solution of the subtitle product of step iii) (11.3g) in DCM (500ml) and stirred for 1h. Saturated sodium thiosulfate solution (100ml) was added and stirred vigorously until no peroxide was detected. The organics were separated and extracted with saturated sodium bicarbonate solution (200ml) and brine (50ml), dried (MgSO_4) and concentrated to yield the subtitle compound as a crude solid. Yield:

25 10.9g.

MS APCI(+ve) 572 $[\text{M}+\text{H}]^+$

v) Sodium 4-((azetidin-1-ylsulfonyl){[2-(trimethylsilyl)ethoxy]methyl}amino)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidine-2-thiolate

30 Sodium hydrosulfide hydrate (1.18g) was added to a solution of the subtitle product of step iv) (8.0g) in DMSO (67ml) and the green solution stirred for 2h. A further aliquot of sodium hydrosulfide hydrate (0.79g) was added and stirred for 1h. This aliquot addition was repeated twice more before heating the solution at 50°C for 30min. The resulting reaction solution was

-76-

used directly in the following step. The subtitle compound was also kept as a stock solution for further reaction with alkyl halides, described in Examples 41-42.

MS APCI(+ve) 450 [M+H]⁺

5 **vi) *N*-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)-*N*-{[2-(trimethylsilyl)ethoxy]methyl}azetidine-1-sulfonamide**


3-Chloro-2-fluorobenzyl bromide (3.13g) was added to an aliquot of the reaction solution of step v) (18ml) containing the subtitle product of step v) and the reaction stirred for 1h. The reaction was partitioned between EtOAc (20ml) and H₂O (20ml) and the organics

10 concentrated *in vacuo*. The residue was purified by column chromatography (20% then 40% EtOAc / iso-hexane) to afford the subtitle compound as an oil which was used directly in the subsequent step.

MS APCI(+ve) 592 [M+H]⁺

15 **Example 41**

***N*-{6-[[*(1R)*-2-Hydroxy-1-methylethyl]amino]-2-[(2,3,4-trifluorobenzyl)thio]-pyrimidin-4-yl}azetidine-1-sulfonamide**



20

A solution of the subtitle product of step i) in trifluoroacetic acid (5ml) was stirred for 15min before the addition of 2M sodium hydroxide solution until pH >10. The aqueous was then extracted with Et₂O (20ml) followed by EtOAc (2 x 20ml). The EtOAc extracts were combined, dried (MgSO₄) and concentrated. The crude material was purified by reverse
25 phase HPLC (90% to 5% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 51mg.

MS APCI(+ve) 464 [M+H]⁺

¹H NMR δ_(DMSO) 10.50 (1H, br. s), 7.47 (1H, m), 7.25 (2H, m), 5.94 (1H, m), 4.70 (1H, br. s), 4.36 (2H, s), 4.04 (1H, br. s), 3.86 (4H, m), 3.40 – 3.25 (2H, m), 2.10 (2H, m), 1.05 (3H, d).

The intermediates for this compound were prepared as follows:


i) *N*-{6-[[*(1R)*-2-Hydroxy-1-methylethyl]amino]-2-[(2,3,4-trifluorobenzyl)thio]-pyrimidin-4-yl}-*N*-{[2-(trimethylsilyl)ethoxy]methyl}azetidine-1-sulfonamide

- 5 2,3,4-Trifluorobenzyl bromide (3.15g) was added to an aliquot of the reaction solution of step v) (18ml) containing the subtitle product of step v) and the reaction stirred for 1h. The reaction was partitioned between EtOAc (20ml) and H₂O (20ml) and the organics concentrated *in vacuo*. The residue was purified by column chromatography (20% then 40% EtOAc / iso-hexane) to afford the subtitle compound as an oil which was used directly in the
- 10 subsequent step.

MS APCI(+ve) 594 [M+H]⁺

Example 42

- N*-(2-[(2,3-Difluoro-4-methylbenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)azetidine-1-sulfonamide
- 15



- A solution of the subtitle product of step i) in trifluoroacetic acid (5ml) was stirred for 15min
- 20 before the addition of 2M sodium hydroxide solution until pH >10. The aqueous was then extracted with Et₂O (20ml) followed by EtOAc (2 x 20ml). The EtOAc extracts were combined, dried (MgSO₄) and concentrated. The crude material was purified by reverse phase HPLC (90% to 5% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 42mg.

- 25 MS APCI(+ve) 460 [M+H]⁺

¹H NMR δ_(DMSO) 7.29 (1H, t), 7.02 (1H, t), 5.93 (1H, s), 4.70 (1H, t), 4.34 (2H, s), 4.05 (1H, br. s), 3.85 (4H, m), 3.41 (1H, m), 3.28 (1H, m), 2.25 (3H, s), 2.10 (2H, quin.), 1.06 (3H, d).

The intermediates for this compound were prepared as follows:

-78-


i) *N*-(2-[(2,3-Difluoro-4-methylbenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)-*N*-{[2-(trimethylsilyl)ethoxy]methyl}azetidine-1-sulfonamide

2,3-Difluoro-4-methylbenzyl bromide (3.10g) was added to an aliquot of the reaction solution of step v) (18ml) containing the subtitle product of step v) and the reaction stirred for 1h. The reaction was partitioned between EtOAc (20ml) and H₂O (20ml) and the organics concentrated *in vacuo*. The residue was purified by column chromatography (20% then 40% EtOAc / iso-hexane) to afford the subtitle compound as an oil which was used directly in the subsequent step.

MS APCI(+ve) 590 [M+H]⁺

Example 43

N-(2-[(2-Fluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)methanesulfonamide



The title product was prepared from a solution of the product of step iv) (4ml) and quenching with 2-fluorobenzyl bromide (0.5g) at room temperature. After 30min the reaction mixture was partitioned between EtOAc (20ml) and saturated brine (20ml). The organic layer was collected and the solvent evaporated to dryness. The residue was passed through a pad of silica gel (1:1 EtOAc / iso-hexane) to give the intermediate product as a gum. This was dissolved in acetonitrile (10ml) and treated with 1M hydrochloric acid with stirring at ambient temperature over 24h. The volatiles were removed *in vacuo* and the residue purified by mass directed reverse phase HPLC to give the title compound as a white foam. Yield: 5mg.

MS APCI(+ve) 387 [M+H]⁺

The intermediates for this compound were prepared as follows:

i) *N*-[2-(benzylthio)-6-chloropyrimidin-4-yl]-*N*-{[2-(trimethylsilyl)ethoxy]methyl}-methanesulfonamide

-79-

To a solution of methanesulfonamide (4.63g) in dry DMF (60ml) at 0°C under nitrogen was added 60% sodium hydride (3.9g) portionwise over 5min. After complete addition the ice bath was removed for 15min and then returned. A solution of the subtitle product of Example 19 step ii) in DMF (30ml) was added dropwise over 5min. After complete addition the ice bath was removed and the whole stirred at ambient temperature for 3h. The ice bath was returned before adding trimethylsilylethoxymethyl chloride (8.6ml) to the reaction mixture. The reaction mixture was allowed to reach ambient temperature and further stirred for 24h. The reaction mixture was partitioned between EtOAc and H₂O. The organic phase collected, dried (MgSO₄) and solvent removed *in vacuo* to leave the subtitle compound as a pale yellow oil. Yield: 21.2g.

MS APCI(+ve) 460 [M+H]⁺

ii) *N*-(2-(Benzylthio)-6-[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-*N*-{[2-(trimethylsilyl)ethoxy]methyl}methanesulfonamide

To a solution of the product of step i) (21.2g) in NMP (40ml) was added (*R*)-alaninol (10g) and the whole heated at 80°C for 4h. The reaction mixture was then partitioned between EtOAc and H₂O. The organic phase was collected, dried (MgSO₄) and the solvent removed to leave a pale yellow oil. This was purified by silica gel chromatography (60:40 iso-hexane / EtOAc) to afford the subtitle compound as a colourless oil. Yield: 13.9g.

MS APCI(+ve) 499 [M+H]⁺

iii) *N*-(2-(Benzylsulfonyl)-6-[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-*N*-{[2-(trimethylsilyl)ethoxy]methyl}methanesulfonamide

To a solution of the product of step ii) (6.8g) in DCM (300ml) was added *m*-chloroperbenzoic acid (8g) at ambient temperature with stirring. After 6h a concentrated solution of sodium thiosulphate (50ml) was added and the organic phase collected. The organic phase was then washed with saturated sodium bicarbonate solution (x2) followed by brine. The organic phase was dried (MgSO₄) and the solvent evaporated to leave the subtitle compound as a colourless foam. Yield: 7.0g.

MS APCI(+ve) 531 [M+H]⁺, MS APCI(-ve) 529 [M-H]⁺

iv) Sodium 4-[(1*R*)-2-hydroxy-1-methylethyl]amino}-6-((methylsulfonyl){[2-(trimethylsilyl)ethoxy]methyl}amino)pyrimidine-2-thiolate


-80-

To a solution of the product of step iii) (3.97g) in DMSO (30ml) was added sodium hydrosulphide hydrate (0.84g). Stirring was continued for a further 2h and additional aliquots of 0.42g of sodium hydrosulphide were added until complete disappearance of starting material as assessed by reverse phase HPLC-MS. The subtitle compound was also kept as a stock solution for further reaction with alkyl halides described in examples 43-53 and 137.

MS APCI(-ve) 407 $[M-H]^+$

Example 44

N-(2-[(2,5-Difluorobenzyl)thio]-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)methanesulfonamide




The title compound was prepared from a solution of the product of Example 43 step iv) (4ml) and quenching with 2,5-difluorobenzyl bromide (0.5g) using the method described for Example 43 to give the title compound as a white foam. Yield: 5mg.

MS APCI(+ve) 405 $[M+H]^+$, MS APCI(-ve) 403 $[M-H]^+$

Example 45

N-(2-[(2,4-Difluorobenzyl)thio]-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)methanesulfonamide



-81-


The title compound was prepared from a solution of the product of Example 43 step iv) (4ml) and quenching with 2,4-difluorobenzyl bromide (0.5g) using the method described for Example 43 to give the title compound as a white foam. Yield: 12mg.

MS APCI(+ve) 405 $[M+H]^+$, MS APCI(-ve) 403 $[M-H]^+$

5

Example 46

N-(2-[(2,6-Difluorobenzyl)thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)methanesulfonamide



10


The title compound was prepared from a solution of the product of Example 43 step iv) (4ml) and quenching with 2,6-difluorobenzyl bromide (0.5g) using the method described for Example 43 to give the title compound as a white foam. Yield: 10mg.

MS APCI(+ve) 405 $[M+H]^+$, MS APCI(-ve) 403 $[M-H]^+$

15

Example 47

N-(2-[(2,3,6-Trifluorobenzyl)thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino}-pyrimidin-4-yl)methanesulfonamide



20


The title compound was prepared from a solution of the product of Example 43 step iv) (4ml) and quenching with 2,3,6-trifluorobenzyl bromide (0.5g) using the method described for Example 43 to give the title compound as a white foam. Yield: 14mg.

-82-

MS APCI(+ve) 423 $[M+H]^+$, MS APCI(-ve) 421 $[M-H]^+$

Example 48


N-(2-[(5-Chloro-2-fluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-
5 pyrimidin-4-yl)methanesulfonamide



The title compound was prepared from a solution of the product of Example 43 step iv) (4ml)
10 and quenching with 5-chloro-2-fluorobenzyl bromide (0.5g) using the method described for
Example 43 to give the title compound as a white foam. Yield: 8mg.
MS APCI(+ve) 421 $[M+H]^+$, MS APCI(-ve) 419 $[M-H]^+$

Example 49

15 *N*-{6-[[*(1R)*-2-Hydroxy-1-methylethyl]amino]-2-[(2,4,5-trifluorobenzyl)thio]-pyrimidin-
4-yl}methanesulfonamide




20 The title compound was prepared from a solution of the product of Example 43 step iv) (4ml)
and quenching with 2,4,5-trifluorobenzyl bromide (0.5g) using the method described for
Example 43 to give the title compound as a white foam. Yield: 7mg.
MS APCI(+ve) 423 $[M+H]^+$, MS APCI(-ve) 421 $[M-H]^+$

Example 50

N-(2-[(3-Chloro-2,6-difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)methanesulfonamide

5




The title compound was prepared from a solution of the product of Example 43 step iv) (4ml) and quenching with 3-chloro-2,6-difluorobenzyl bromide (0.5g) using the method described for Example 43 to give the title compound as a white foam. Yield: 27mg.

MS APCI(+ve) 439 [M+H]⁺

Example 51

N-{6-[[*(1R)*-2-Hydroxy-1-methylethyl]amino]-2-[(2,4,6-trifluorobenzyl)thio]-pyrimidin-4-yl}methanesulfonamide




The title compound was prepared from a solution of the product of Example 43 step iv) (4ml) and quenching with 2,4,6-trifluorobenzyl bromide (0.5g) using the method described for Example 43 to give the title compound as a white foam. Yield: 42mg.

MS APCI(+ve) 423 [M+H]⁺

Example 52

***N*-(2-[(2-Chloro-3,6-difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)methanesulfonamide**

5



The title compound was prepared from a solution of the product of Example 43 step iv) (4ml) and quenching with 2-chloro-3,6-difluorobenzyl bromide (0.5g) using the method described


10 for Example 43 to give the title compound as a white foam. Yield: 40mg.

MS APCI(+ve) 439 [M+H]⁺

Example 53

***N*-(2-[(2-Chloro-6-fluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-**

15 **pyrimidin-4-yl)methanesulfonamide**



The title compound was prepared from a solution of the product of Example 43 step iv) (4ml)

20 and quenching with 2-chloro-6-fluorobenzyl bromide (0.5g) using the method described for Example 43 to give the title compound as a white foam. Yield: 32mg.

MS APCI(+ve) 421 [M+H]⁺


General Procedure for the Synthesis of Examples 54 to 99.

To the required sulfonyl chloride (0.15mM) was added a solution of the subtitle product of Example 3 step ii) (0.05mM) in pyridine (0.4ml) and 4-*N,N*-dimethylaminopyridine (0.05mM) in pyridine (0.2ml) before the reaction mixture was stirred at room temperature for three days. 3M Hydrochloric acid (0.2ml) was added and stirring maintained for 18h before the solvent was removed under reduced pressure. The residue was dissolved in DMSO / H₂O (400μl; 3:1), filtered through a PORVAIR filter and the product purified by mass directed reverse phase HPLC to afford the title products of Examples 54 to 99.

10

Example 54

***N*-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-4-methylbenzenesulfonamide**




Yield: 7mg.

MS: APCI (+ve) 369 [M+H]⁺

20 **Example 55**

***N*-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-2,4,6-trimethylbenzenesulfonamide.**




-86-

Yield: 11mg.

MS: APCI (+ve) 473[M+H]⁺**5 Example 56**

N-{2-(Benzylthio)-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl}naphthalene-2-sulfonamide.




10

Yield: 14mg.

MS: APCI (+ve) 481 [M+H]⁺**Example 57**

- 15 *N*-(2-(Benzylthio)-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)-1-(*R,S*)-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethanesulfonamide



20 Yield: 4mg.


MS: APCI (+ve) 528 [M+H]⁺

25

-87-

Example 58

***N*-{2-(Benzylthio)-6-[[[(1*R*)-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl}-4-bromobenzenesulfonamide**




Yield: 7mg.

MS: APCI (+ve) 509/511 $[M+H]^+$

10 **Example 59**

***N*-{2-(Benzylthio)-6-[[[(1*R*)-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl}-4-tert-butylbenzenesulfonamide**



15

Yield: 14mg.


MS: APCI (+ve) 487 $[M+H]^+$

20

-88-

Example 60


***N*-{2-(Benzylthio)-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl}-2-bromobenzenesulfonamide**



MS: APCI (+ve) 509/511 $[M+H]^+$

Example 61

10 ***N*-{2-(Benzylthio)-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl}-4-(trifluoromethyl)benzenesulfonamide**




15 Yield: 14mg.

MS: APCI (+ve) 499 $[M+H]^+$

Example 62

20 ***N*-{2-(Benzylthio)-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl}-3-(trifluoromethyl)benzenesulfonamide**

-89-




Yield: 9mg.

MS: APCI (+ve) 499 $[M+H]^+$

5

Example 63

***N*-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-2,5-dimethoxybenzenesulfonamide**




10

Yield: 13mg.

MS: APCI (+ve) 491 $[M+H]^+$ **Example 64**

***N*-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-2,1,3-benzoxadiazole-4-sulfonamide**

-90-




Yield: 4mg.

MS: APCI (+ve) 473 [M+H]⁺

5

Example 65

N-{2-(Benzylthio)-6-(((1*R*)-2-hydroxy-1-methylethyl)amino)pyrimidin-4-yl}-5-isoxazol-3-ylthiophene-2-sulfonamide




10

Yield: 8mg.

MS: APCI (+ve) 504 [M+H]⁺15 **Example 66**

N-{2-(Benzylthio)-6-(((1*R*)-2-hydroxy-1-methylethyl)amino)pyrimidin-4-yl}-2,6-dichlorobenzenesulfonamide




20

-91-

Yield: 10mg.

MS: APCI (+ve) 499/501/503 $[M+H]^+$ **Example 67**


- 5 *N*-{2-(Benzylthio)-6-[[[(1*R*)-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl]-2,6-difluorobenzenesulfonamide



- 10 Yield: 7mg.

MS: APCI (+ve) 467 $[M+H]^+$ **Example 68**

- 15 *N*-{2-(Benzylthio)-6-[[[(1*R*)-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl]-4-(1,1-dimethylpropyl)benzenesulfonamide




Yield: 13mg.

- 20 MS: APCI (+ve) 501 $[M+H]^+$

Example 69

- N*-{2-(Benzylthio)-6-[[[(1*R*)-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl]-2-chloro-4-fluorobenzenesulfonamide

-92-




Yield: 9mg.

5 MS: APCI (+ve) 483/485 [M+H]⁺**Example 70**

***N*-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-3-chloro-4-fluorobenzenesulfonamide**

10



Yield: 15mg.


MS: APCI (+ve) 483/485 [M+H]⁺

15

Example 71

***N*-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-2,5-dichlorobenzenesulfonamide**

-93-




Yield: 11mg.

MS: APCI (+ve) 499/501/503 [M+H]⁺

5

Example 72


N-{2-(Benzylthio)-6-(((1*R*)-2-hydroxy-1-methylethyl)amino)pyrimidin-4-yl}-4-propylbenzenesulfonamide



Yield 14mg.

MS: APCI (+ve) 473 [M+H]⁺**Example 73**

N-{2-(Benzylthio)-6-(((1*R*)-2-hydroxy-1-methylethyl)amino)pyrimidin-4-yl}-3-bromobenzenesulfonamide




20

-94-

Yield: 17mg.

MS: APCI (+ve) 509/511 $[M+H]^+$ **Example 74**


- 5 *N*-{2-(Benzylthio)-6-[(*(1R)*-2-hydroxy-1-methylethyl)amino]pyrimidin-4-yl}-3-chloro-2-methylbenzenesulfonamide



- 10 Yield: 13mg.

MS: APCI (+ve) 479/481 $[M+H]^+$ **Example 75**

- N*-{2-(Benzylthio)-6-[(*(1R)*-2-hydroxy-1-methylethyl)amino]pyrimidin-4-yl}-2,5-dichlorothiophene-3-sulfonamide




Yield: 6mg.

- 20 MS: APCI (+ve) 505/507/509 $[M+H]^+$

Example 76

- N*-{2-(Benzylthio)-6-[(*(1R)*-2-hydroxy-1-methylethyl)amino]pyrimidin-4-yl}-3,4-dimethoxybenzenesulfonamide

-95-




Yield: 11mg.

5 MS: APCI (+ve) 491 $[M+H]^+$ **Example 77**

N-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-2,3-dichlorobenzenesulfonamide

10



Yield: 11mg.


MS: APCI (+ve) 499/501/503 $[M+H]^+$

15

Example 78

N-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-5-chlorothiophene-2-sulfonamide

-96-




Yield: 8mg.

MS: APCI (+ve) 471/473 [M+H]⁺

5

Example 79

***N*-{2-(Benzylthio)-6-(((1*R*)-2-hydroxy-1-methylethyl)amino)pyrimidin-4-yl}-2-chloro-6-methylbenzenesulfonamide**




10

Yield: 7mg.

MS: APCI (+ve) 479/481 [M+H]⁺**15 Example 80**

***N*-{2-(Benzylthio)-6-(((1*R*)-2-hydroxy-1-methylethyl)amino)pyrimidin-4-yl}-3,4-dichlorobenzenesulfonamide**




Yield: 13mg.

MS: APCI (+ve) 499/501/503 $[M+H]^+$

5 **Example 81**

***N*-{2-(Benzylthio)-6-[(*(1R)*-2-hydroxy-1-methylethyl)amino]pyrimidin-4-yl}-3,5-dichlorobenzenesulfonamide**




10

Yield: 14mg.

MS: APCI (+ve) 499/501/503 $[M+H]^+$

Example 82

15 ***N*-{2-(Benzylthio)-6-[(*(1R)*-2-hydroxy-1-methylethyl)amino]pyrimidin-4-yl}-2,4-dichloro-5-methylbenzenesulfonamide**




20 Yield: 12mg.

MS: APCI (+ve) 513/515/517 $[M+H]^+$

Example 83

N-{2-(Benzylthio)-6-[(*(1R)*-2-hydroxy-1-methylethyl)amino]pyrimidin-4-yl}-3,4-difluorobenzenesulfonamide




Yield: 15mg.

MS: APCI (+ve) 467 [M+H]⁺

10 **Example 84**

N-{2-(Benzylthio)-6-[(*(1R)*-2-hydroxy-1-methylethyl)amino]pyrimidin-4-yl}-2-methylbenzenesulfonamide



15


Yield: 14mg.

MS: APCI (+ve) 445 [M+H]⁺

Example 85

20 *N*-{2-(Benzylthio)-6-[(*(1R)*-2-hydroxy-1-methylethyl)amino]pyrimidin-4-yl}-3-methoxybenzenesulfonamide

-99-




Yield: 16mg.

MS: APCI (+ve) 461 [M+H]⁺

5

Example 86

***N*-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-2,5-difluorobenzenesulfonamide**




10

Yield: 12mg.

MS: APCI (+ve) 467 [M+H]⁺15 **Example 87**

***N*-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-4-chloro-2,5-dimethoxybenzenesulfonamide**




20

-100-

Yield: 9mg.

MS: APCI (+ve) 525/527 $[M+H]^+$ **Example 88**


- 5 *N*-{2-(Benzylthio)-6-(((1*R*)-2-hydroxy-1-methylethyl)amino)pyrimidin-4-yl}thiophene-3-sulfonamide



- 10 Yield: 18mg.

MS: APCI (+ve) 437 $[M+H]^+$ **Example 89**

- 15 4-Acetyl-*N*-{2-(benzylthio)-6-(((1*R*)-2-hydroxy-1-methylethyl)amino)pyrimidin-4-yl}benzenesulfonamide




Yield: 6mg.

- 20 MS: APCI (+ve) 473 $[M+H]^+$

Example 90

- N*-{2-(Benzylthio)-6-(((1*R*)-2-hydroxy-1-methylethyl)amino)pyrimidin-4-yl}-2-chloro-4,5-difluorobenzenesulfonamide




Yield: 5mg.

5 MS: APCI (+ve) 501/503 $[M+H]^+$

Example 91

N-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-5-chloro-2,4-difluorobenzenesulfonamide

10



Yield: 4mg.


MS: APCI (+ve) 501/503 $[M+H]^+$

15

Example 92

N-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-4-chloro-2,5-difluorobenzenesulfonamide

-102-




Yield: 9mg.

MS: APCI (+ve) 501/503 $[M+H]^+$

5

Example 93

N-{2-(Benzylthio)-6-[(*(1R)*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}biphenyl-4-sulfonamide




10

Yield: 14mg.

MS: APCI (+ve) 507 $[M+H]^+$ **Example 94**

N-{2-(Benzylthio)-6-[(*(1R)*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-2-methoxy-4-methylbenzenesulfonamide

-103-




Yield: 10mg.

MS: APCI (+ve) 475 $[M+H]^+$

5

Example 95

***N*-{2-(Benzylthio)-6-[(*(1R)*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-3-chloro-4-methylbenzenesulfonamide**




10

Yield: 14mg.

MS: APCI (+ve) 479/481 $[M+H]^+$ 15 **Example 96**

***N*-{2-(Benzylthio)-6-[(*(1R)*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-4-bromo-2-methylbenzenesulfonamide**




-104-

Yield: 5mg.

MS: APCI (+ve) 523/525 $[M+H]^+$ **5 Example 97**

***N*-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-4-phenoxybenzenesulfonamide**




10

Yield: 7 mg.

MS: APCI (+ve) 523 $[M+H]^+$ **Example 98**

***N*-{2-(Benzylthio)-6-[(*1R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl}-4-chloro-2,5-dimethylbenzenesulfonamide**




20 Yield: 14mg.

MS: APCI (+ve) 493/495 $[M+H]^+$

Example 99

***N*-{2-(Benzylthio)-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl}-2,3,4-trifluorobenzenesulfonamide**



Yield: 5mg.

MS: APCI (+ve) 485 [M+H]⁺


10 **General Procedure for the Synthesis of Examples 100 to 105.**

To the required sulfonyl chloride (0.15mM) was added a solution of the subtitle product of Example 3 step ii) (0.05mM) in pyridine (0.4ml) and 4-*N,N*-dimethylaminopyridine (0.05mM) in pyridine (0.2ml) before the reaction mixture was stirred at room temperature for three days. 3M Hydrochloric acid (0.2ml) was added and stirring maintained for 18h before
15 the solvent was removed under reduced pressure. The residue was dissolved in DMSO / H₂O (400μl; 3:1) and filtered through a PORVAIR filter before the product was purified by mass directed reverse phase HPLC to afford the title products of Examples 100 to 105 as solution samples.


20 **Example 100**

***N*-(2-(Benzylthio)-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)-4-methoxybenzenesulfonamide**

-106-

MS: APCI (+ve) 461 $[M+H]^+$ **5 Example 101**

N-(2-(Benzylthio)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-2,4-dichlorobenzenesulfonamide




10

MS: APCI (+ve) 499 $[M+H]^+$ **Example 102**

N-(2-(Benzylthio)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)thiophene-2-sulfonamide

15


MS: APCI (+ve) 437 $[M+H]^+$

20

-107-

Example 103


***N*-(2-(Benzylthio)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-2,5-dimethylbenzenesulfonamide**



MS: APCI (+ve) 459 $[M+H]^+$

Example 104

10 ***N*-(2-(Benzylthio)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-1,2-dimethyl-1*H*-imidazole-4-sulfonamide**




15 MS: APCI (+ve) 449 $[M+H]^+$

Example 105

***N*-(2-(Benzylthio)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)butane-1-sulfonamide**

20


-108-



MS: APCI (+ve) 411 $[M+H]^+$

5 Example 106

N-(2-[(2,3-Difluorobenzyl)thio]-6-[[*(1R)*]-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)morpholine-4-sulfonamide



10

2,3-Difluorobenzyl bromide (0.95g) was added to an aliquot of the reaction solution of step iv) (2ml) containing the subtitle product of step iv) and the reaction stirred for 2h. The reaction was partitioned between EtOAc (20ml) and brine (20ml). The aqueous was extracted with EtOAc (2 x 20ml) and the organics concentrated *in vacuo*. The residue was purified by

15 column chromatography (30% then 40% EtOAc / iso-hexane) to afford the subtitle product as an oil that was diluted in acetonitrile (5ml) and 2M hydrochloric acid (5ml) and was stirred overnight before removal of the volatiles *in vacuo*. The crude material was purified by reverse phase HPLC (gradient 95% to 20% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 0.17g.

20 MS APCI(+ve) 476 $[M+H]^+$

^1H NMR $\delta_{(\text{DMSO})}$ 10.57 (1H, bs), 7.40 (1H, bt), 7.32 (2H, m), 7.15 (1H, m), 5.90 (1H, s), 4.71 (1H, bs), 4.39 (2H, t), 4.02 (1H, bs), 3.60 (4H, t), 3.40 (1H, m), 3.30 (1H, m), 3.18 (4H, bs), 1.06 (3H, d).

-109-

The intermediates for this compound were prepared as follows:

i) *N*-{6-Chloro-2-[benzylthio]pyrimidin-4-yl}-*N*-{[2-(trimethylsilyl)ethoxy]methyl}morpholine-4-sulfonamide

The subtitle compound was prepared as an oil by the method of Example 32 step i) (8.9g)

5 using the subtitle product of Example 19 step ii) and the subtitle product of Example 36 step i) (4.7g) and 2-(trimethylsilyl)ethoxymethyl chloride (6.1g). Yield: 11.8g.

MS APCI(+ve) 401 [M+H]⁺

10 ii) *N*-(2-[benzylthio]-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}-pyrimidin-4-yl)-*N*-{[2-(trimethylsilyl)ethoxy]methyl}morpholine-4-sulfonamide

The subtitle compound was prepared as a yellow oil by the method of Example 43 step ii) by reacting the subtitle product of step i) (11.75g) with (*R*)-alaninol (3.4ml) in NMP (30ml).

Yield: 12.2g.

MS APCI(+ve) 570 [M+H]⁺

15

iii) *N*-(2-[benzylsulfonyl]-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}-pyrimidin-4-yl)-*N*-{[2-(trimethylsilyl)ethoxy]methyl}morpholine-4-sulfonamide

m-Chloroperbenzoic acid (5.87g) was added as a single portion to a solution of the subtitle product of step iii) (5.83g) in DCM (220ml) and stirred for 2.5h. A further aliquot of *m*-

20 chloroperbenzoic acid (1.0g) was added and stirring maintained for 1h. Saturated sodium thiosulfate solution (100ml) was added and stirred vigorously until no peroxides were detected. The organics were separated and extracted with saturated sodium bicarbonate solution (200ml) and brine (50ml), dried (MgSO₄) and concentrated to yield the subtitle compound as a crude beige white solid. Yield: 5.6g.

25 MS APCI(+ve) 602 [M+H]⁺

iv) Sodium 4-{[(1*R*)-2-hydroxy-1-methylethyl]amino}-6-((morpholin-4-ylsulfonyl){[2-(trimethylsilyl)ethoxy]methyl}amino)pyrimidine-2-thiolate

Sodium hydrosulfide hydrate (0.62g) was added to a solution of the subtitle product of step

30 iii) (2.5g) in DMSO (5ml) and the green solution stirred for 1.25h. A further aliquot of sodium hydrosulfide hydrate (0.28g) was added and stirred for 45min. A further aliquot of sodium hydrosulfide hydrate (0.32g) was added and stirred for 1.25h before the addition of a final aliquot of sodium hydrosulfide hydrate (0.10g) in DMSO (1ml). The resulting reaction

-110-


solution was diluted with DMSO (10ml) and used directly in the following step. The subtitle compound was also kept as a stock solution for further reaction with alkyl halides, described in Examples 107-110.

MS APCI(+ve) 480 [M+H]⁺

5

Example 107

N-{6-{[(1*R*)-2-Hydroxy-1-methylethyl]amino}-2-[(2,3,4-trifluorobenzyl)thio]-pyrimidin-4-yl}morpholine-4-sulfonamide



2,3,4-Trifluorobenzyl bromide (1.04g) was added to an aliquot of the reaction solution of Example 106 step iv) (2ml) containing the subtitle product of Example 106 step iv) and the reaction stirred for 2h. The reaction was partitioned between EtOAc (20ml) and brine (20ml).

15 The aqueous was extracted with EtOAc (2 x 20ml) and the organics concentrated *in vacuo*. The residue was purified by column chromatography (50% then 60% EtOAc / iso-hexane) to afford the subtitle product as an oil that was diluted in acetonitrile (5ml) and 2M hydrochloric acid (5ml) and was stirred overnight before removal of the volatiles *in vacuo*. The crude material was purified by reverse phase HPLC (gradient 95% to 20% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 0.14g.


MS APCI(+ve) 494 [M+H]⁺

¹H NMR δ(DMSO) 10.57 (1H, bs), 7.45 (1H, bs), 7.25 (2H, m), 5.90 (1H, s), 4.71 (1H, bs), 4.36 (2H, s), 4.02 (1H, s), 3.60 (4H, bs), 3.38 (1H, m), 3.30 (1H, m), 3.15 (4H, bs), 1.07 (3H, d).

25 **Example 108**

N-{6-{[(1*R*)-2-Hydroxy-1-methylethyl]amino}-2-[(2,3,5-trifluorobenzyl)thio]-pyrimidin-4-yl}morpholine-4-sulfonamide

-111-




2,3,5-Trifluorobenzyl bromide (1.04g) was added to an aliquot of the reaction solution of Example 106 step iv) (2ml) containing the subtitle product of Example 106 step iv) and the reaction stirred for 2h. The reaction was partitioned between EtOAc (20ml) and brine (20ml). The aqueous was extracted with EtOAc (2 x 20ml) and the organics concentrated *in vacuo*. The residue was purified by column chromatography (50% then 60% EtOAc / iso-hexane) to afford the subtitle product as an oil that was diluted in acetonitrile (5ml) and 2M hydrochloric acid (5ml) and was stirred overnight before removal of the volatiles *in vacuo*. The crude material was purified by reverse phase HPLC (gradient 95% to 20% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 0.16g.

MS APCI(-ve) 492 [M-H]⁻

¹H NMR $\delta_{(\text{DMSO})}$ 10.61 (1H, bs), 7.38 (3H, bm), 5.91 (1H, s), 4.71 (1H, bs), 4.36 (2H, t), 4.02 (1H, bs), 3.59 (4H, bs), 3.37 (1H, m), 3.30 (1H, m), 3.14 (4H, bs), 1.05 (3H, d).

Example 109

***N*-(2-[(2,3-Difluoro-4-methylbenzyl)thio]-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)morpholine-4-sulfonamide**



2,3-Difluoro-4-methylbenzyl bromide (1.02g) was added to an aliquot of the reaction solution of Example 106 step iv) (2ml) containing the subtitle product of Example 106 step iv) and the reaction stirred for 2h. The reaction was partitioned between EtOAc (20ml) and

-112-


brine (20ml). The aqueous was extracted with EtOAc (2 x 20ml) and the organics concentrated *in vacuo*. The residue was purified by column chromatography (70% EtOAc / iso-hexane) to afford the subtitle product as an oil that was diluted in acetonitrile (5ml) and 2M hydrochloric acid (5ml) and was stirred overnight before removal of the volatiles *in vacuo*. The crude material was purified by reverse phase HPLC (gradient 95% to 20% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 0.12g.

MS APCI(-ve) 488 [M-H]⁻

¹H NMR $\delta_{(\text{DMSO})}$ 10.57 (1H, bs), 7.28 (2H, bs), 7.02 (1H, t), 5.90 (1H, bs), 4.71 (1H, bs), 4.34 (2H, bm), 4.03 (1H, bs), 3.59 (4H, bs), 3.39 (1H, m), 3.30 (1H, m), 3.15 (4H, bs), 2.24 (3H, s), 1.07 (3H, d).

Example 110

N-(2-[(4-Chloro-2-fluorobenzyl)thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino)-pyrimidin-4-yl)morpholine-4-sulfonamide



4-Chloro-2-fluorobenzyl bromide (1.03g) was added to an aliquot of the reaction solution of Example 106 step iv) (2ml) containing the subtitle product of Example 106 step iv) and the reaction stirred for 2h. The reaction was partitioned between EtOAc (20ml) and brine (20ml). The aqueous was extracted with EtOAc (2 x 20ml) and the organics concentrated *in vacuo*. The residue was purified by column chromatography (66% EtOAc / iso-hexane) to afford the subtitle product as an oil that was diluted in acetonitrile (5ml) and 2M hydrochloric acid (5ml) and was stirred overnight before removal of the volatiles *in vacuo*. The crude material was purified by reverse phase HPLC (gradient 95% to 20% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 90mg.


MS APCI(-ve) 490 [M-H]⁻

-113-

¹H NMR $\delta_{(\text{DMSO})}$ 10.56 (1H, bs), 7.63 (1H, bt), 7.42 (1H, d), 7.31 (1H, bs), 7.23 (1H, d), 5.90 (1H, s), 4.72 (1H, bs), 4.32 (2H, bs), 4.30 (1H, bs), 3.59 (4H, bs), 3.40 (1H, m), 3.30 (1H, m), 3.15 (4H, bs), 1.06 (3H, d).

5 Example 111

N-(2-[(2,3-Difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)pyrrolidine-1-sulfonamide



10

2,3-Difluorobenzyl bromide (2.65g) was added to an aliquot of the reaction solution of step v) (12.6ml) containing the subtitle product of step v) and the reaction stirred for 1h. The reaction was partitioned between EtOAc (20ml) and H₂O (20ml), the organics were recovered, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography (650:350:1 iso-hexane / EtOAc / AcOH) to afford the subtitle compound as an oil that was diluted in trifluoroacetic acid (2ml) and was stirred for 12min before quenching the reaction by the addition of 1M sodium hydroxide solution to pH >10. The aqueous was washed with Et₂O before saturated ammonium chloride solution was added to acidify the aqueous to pH 4 followed by extracting with EtOAc (3 x 20ml). The EtOAc extracts were combined, dried (MgSO₄) and concentrated *in vacuo*. The crude material was purified by reverse phase HPLC (gradient 95% to 20% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 0.52g.

MS APCI(+ve) 460 [M+H]⁺

¹H NMR $\delta_{(\text{CDCl}_3)}$ 7.25 - 7.20 (1H, m), 7.08 - 6.97 (2H, m), 5.95 (1H, s), 4.98 (1H, d), 4.34 (2H, s), 4.15 - 4.01 (1H, m), 3.73 - 3.69 (1H, m), 3.60 - 3.55 (1H, m), 3.39 (4H, t), 1.93 - 1.90 (4H, m), 1.21 (3H, d).

The intermediates for this compound were prepared as follows:

i) Pyrrolidine-1-sulfonamide

-114-

Pyrrolidine (3.37g) and sulfamide (7.10g) in 1,4-dioxane (110ml) were heated at reflux for 24h. The solvent was evaporated under reduced pressure and the resulting solid suspended in CHCl_3 . The suspension was filtered and the filtrate concentrated *in vacuo* to afford the subtitle compound as a white solid. Yield: 5.35g.

5 ^1H NMR $\delta_{(\text{CDCl}_3)}$ 4.46 (2H, s), 3.31 (4H, t), 1.96-1.92 (4H, m).

ii) *N*-[2-(Benzylthio)-6-chloropyrimidin-4-yl]-*N*-{[2-(trimethylsilyl)ethoxy]methyl}-pyrrolidine-1-sulfonamide

To a solution of the product of step i) (5.0g) in dry DMF (60ml) at 0°C under nitrogen was
10 added 60% sodium hydride (2.66g). The reaction was allowed to warm outside the cooling bath for 15min before recooling to 0°C and addition of the product from Example 19 step ii) (9.03g) in DMF (20ml) and the whole allowed to further stir at room temperature for 3h. The reaction was quenched with 2-(trimethylsilyl)ethoxymethyl chloride (6.50ml) and allowed to stir for 18h before removal of the volatiles *in vacuo* and partitioning of the residue between
15 EtOAc (100ml) and H_2O (200ml). The aqueous was washed further with EtOAc (2 x 100ml) and the organics combined, dried (MgSO_4) and concentrated *in vacuo*. The residue was purified by column chromatography (1:18:181 AcOH / EtOAc / iso-hexane) to afford the subtitle compound as a colourless oil. Yield: 8.26g.

MS APCI(+ve) 515 $[\text{M}+\text{H}]^+$

20

iii) *N*-(2-(Benzylthio)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-*N*-{[2-(trimethylsilyl)ethoxy]methyl}pyrrolidine-1-sulfonamide

The subtitle compound was prepared as a yellow oil by the method of Example 43 step ii) by reacting the subtitle product of step ii) (8.26g) with (*R*)-alaninol (3.61g) in NMP (60ml).

25 Yield: 7.6g.

MS APCI(+ve) 554 $[\text{M}+\text{H}]^+$

iv) *N*-(2-(Benzylsulfonyl)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-*N*-{[2-(trimethylsilyl)ethoxy]methyl}pyrrolidine-1-sulfonamide

30 *m*-Chloroperbenzoic acid (11.07g) was added as a single portion to a solution of the subtitle product of step iii) (9.41g) in DCM (44ml) and stirred for 6h. Saturated sodium thiosulfate solution (100ml) was added and stirred vigorously until no peroxides were detected. The organics were separated and extracted with saturated sodium bicarbonate solution (200ml)

-115-

and brine (50ml), dried (MgSO₄) and concentrated to yield the subtitle compound as a colourless foam. Yield: 1.0g.

MS APCI(+ve) 531 [M+H]⁺

5 v) **Sodium 4-{[(1*R*)-2-hydroxy-1-methylethyl]amino}-6-((pyrrolidin-1-ylsulfonyl){[2-(trimethylsilyl)ethoxy]methyl}amino)pyrimidine-2-thiolate**


Sodium hydrosulfide hydrate (2.15g) was added to a solution of the subtitle product of step iii) (4.5g) in DMSO (37.8ml) and the green solution stirred for 1h. A further aliquot of sodium hydrosulfide hydrate (0.1g) was added and stirred for 1h. A further aliquot of sodium hydrosulfide hydrate (0.1g) was added and stirred for 2h before the addition of a final aliquot of sodium hydrosulfide hydrate (0.05g). The resulting reaction solution was used directly in the following step. The subtitle compound was also kept as a stock solution for further reaction with alkyl halides, described in Examples 112-113.

MS APCI(+ve) 351 [M+H]⁺

15

Example 112

***N*-{6-{[(1*R*)-2-Hydroxy-1-methylethyl]amino}-2-[(2,3,4-trifluorobenzyl)thio]-pyrimidin-4-yl}pyrrolidine-1-sulfonamide**



20

The title compound was prepared as a white solid by the method of Example 111 using the subtitle product of Example 111 step v) (12.6ml) and 2,3,4-trifluorobenzyl bromide (2.88g). Yield: 0.12g.


25 MS APCI(+ve) 478 [M+H]⁺

¹H NMR δ_(CDCl₃) 7.22 - 7.16 (1H, m), 7.01 - 6.86 (2H, m), 5.95 (1H, s), 5.01 (1H, d), 4.30 (2H, s), 4.07 (1H, m), 3.74 - 3.70 (1H, m), 3.60 - 3.56 (1H, m), 3.39 (4H, t), 1.94 - 1.90 (4H, m), 1.23 (3H, d).

-116-

Example 113

***N*-(2-[(2,3-Difluoro-4-methylbenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)pyrrolidine-1-sulfonamide**




The title compound was prepared as a white solid by the method of Example 111 using the subtitle product of Example 111 step v) (12.6ml) and 2,3-difluoro-4-methylbenzyl bromide (2.83g). Yield: 40mg.

10 MS APCI(-ve) 372 [M-H]⁻

¹H NMR δ_(DMSO) 10.36 (1H, s), 7.27 (1H, t), 7.01 (1H, t), 5.78 (1H, s), 4.69 (1H, t), 4.32 (2H, s), 4.03 - 3.87 (1H, m), 3.33 - 3.29 (1H, m), 3.28 - 3.22 (4H, m), 2.24 (3H, s), 1.78 - 1.75 (4H, m), 1.06 (3H, d).

15 **Example 114**

***N*-(2-[(2,3-Difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)piperidine-1-sulfonamide**



20

The subtitle product of step ii) was heated in (*R*)-alaninol (2ml) for 8 days at 80°C before partitioning between EtOAc (50ml) and H₂O (50ml). The organics were recovered, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography

-117-

(550:450:1 EtOAc / iso-hexane / AcOH) to afford the title compound as a white solid. Yield: 106mg.

MS APCI(+ve) 474 [M+H]⁺

¹H NMR $\delta_{(\text{CDCl}_3)}$ 7.24 - 7.20 (1H, m), 7.08 - 6.98 (2H, m), 6.02 (1H, s), 4.36 (2H, s), 4.18 - 3.96 (1H, m), 3.74 - 3.70 (1H, m), 3.61 - 3.57 (1H, m), 3.26 (4H, t), 1.65 - 1.59 (4H, m), 1.57 - 1.51 (2H, m), 1.22 (3H, d).

The intermediates for this compound were prepared as follows:

i) Piperidine-1-sulfonamide

- 10 Piperidine (3.0g) and sulfamide (5.93g) in 1,4-dioxane (100ml) were heated at reflux for 24h. The solvent was evaporated under reduced pressure and the resulting solid suspended in CHCl₃. The suspension was filtered and the filtrate concentrated *in vacuo* to afford the subtitle compound as a white solid. Yield: 3.85g.

¹H NMR $\delta_{(\text{DMSO})}$ 6.65 (2H, s), 2.92 (4H, t), 1.59-1.53 (4H, m), 1.45-1.40 (2H, m).

15

ii) N-{3-Chloro-5-[(2,3-difluorobenzyl)thio]phenyl}piperidine-1-sulfonamide


- 60% Sodium hydride (0.20g) was added to a solution of the subtitle product of step i) (0.4g) in DMF (6.7ml) at 0°C. The reaction was allowed to warm outside the cooling bath for 15min before recooling to 0°C for 15min. A solution of the subtitle product of Example 39 step ii) (0.75g) in DMF (2ml) was then added and stirring maintained for 3h. The reaction mixture was neutralised with methanolic hydrogen chloride before concentrating *in vacuo*. The residue was partitioned between EtOAc (100ml) and H₂O (200ml) and the organics recovered, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography (1:20:79 AcOH / EtOAc / iso-hexane) to afford the subtitle compound as a colourless oil. Yield: 1.3g.

MS APCI(+ve) 435 [M+H]⁺

Example 115

- N-(2-[(2-Fluoro-3-methylbenzyl)thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino)-pyrimidin-4-yl)methanesulfonamide**
- 30

-118-



A solution of the subtitle product of step iii) (1.0g) in (*R*)-alaninol (1.5ml) was heated at 80°C for 18h before partitioning between EtOAc and H₂O. The organics were recovered, dried (MgSO₄) and concentrated. The residue was purified by column chromatography (1:76:133 AcOH / EtOAc / iso-hexane) before diluting the crude material in trifluoroacetic acid (2ml) and stirring for 12min before quenching the reaction by the addition of 1M sodium hydroxide solution to pH >10. The aqueous was extracted with Et₂O before being acidified with saturated ammonium chloride solution to pH 4 and extracting with EtOAc (3 x 20ml). The EtOAc extracts were dried (MgSO₄) and concentrated *in vacuo*. The crude material was purified by reverse phase HPLC (gradient 95% to 20% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 0.33g.

MS APCI(+ve) 401 [M+H]⁺

¹H NMR δ_(DMSO) 10.53 (1H, s), 7.36 (1H, t), 7.17 (1H, t), 7.01 (1H, t), 5.77 (1H, s), 4.70 (1H, s), 4.33 (2H, s), 4.01 (1H, s), 3.42 - 3.37 (1H, m), 3.32 (3H, s), 3.31 - 3.26 (1H, m), 2.23 (3H, s), 1.07 (3H, d).

The intermediates for this compound were prepared as follows:

i) 2-[(2-Fluoro-3-methylbenzyl)thio]pyrimidine-4,6-diol

The subtitle compound was prepared as a yellow solid by the method of Example 39 step i) using 2-fluoro-3-methylbenzyl bromide (7.0g), 2-mercaptopyrimidine-4,6-diol (5.0g) and potassium hydroxide (1.93g). Yield: 8.36g.

MS APCI(+ve) 267 [M+H]⁺

ii) 4,6-Dichloro-2-[(2-fluoro-3-methylbenzyl)thio]pyrimidine

The subtitle compound was prepared as white crystals by the method of Example 39 step ii) using the subtitle product of step i) (8.36g), phosphorus oxychloride (47ml) and *N,N*-dimethylaniline (8.9ml). Yield: 7.32g.

-119-

^1H NMR $\delta_{(\text{CDCl}_3)}$ 7.32 (1H, t), 7.10 (1H, t), 7.03 (1H, s), 6.96 (1H, t), 4.40 (2H, s), 2.28 (3H, s).


iii) *N*-{6-Chloro-2-[(2-fluoro-3-methylbenzyl)thio]pyrimidin-4-yl}-*N*-{[2-(trimethylsilyl)ethoxy]methyl}azetidine-1-sulfonamide

The subtitle compound was prepared as a colourless oil by the method of Example 39 step iii) using the subtitle product of step ii) (2.45g), methanesulfonamide (0.76g), 60% sodium hydride (0.64g) and 2-(trimethylsilyl)ethoxymethyl chloride (1.42ml). Yield: 3.14g. MS APCI(+ve) 434 $[\text{M}+\text{H}]^+$

10

Example 116

N-{2-[(2-Fluoro-3-methylbenzyl)thio]-6-[(2-hydroxy-1,1-dimethylethyl)amino]-pyrimidin-4-yl}methanesulfonamide



15

The title compound was prepared as a white solid by the method of Example 115 using the subtitle product of Example 115 step iii) (1.0g), 2-amino-2-methyl-1-propanol (1.5ml) and trifluoroacetic acid (2ml). Yield: 0.19g.


20 MS APCI(+ve) 415 $[\text{M}+\text{H}]^+$

^1H NMR $\delta_{(\text{CDCl}_3)}$ 7.25 (1H, t), 7.08 (1H, t), 6.97 (1H, t), 5.95 (1H, s), 4.94 (1H, s), 4.34 (2H, s), 3.64 (2H, s), 3.16 (3H, s), 2.28 (3H, s), 1.36 (6H, s).

Example 117

25 *N*-(2-[(2-Fluoro-3-methylbenzyl)thio]-6-[(1*R*)-1-(hydroxymethyl)propyl]amino)-pyrimidin-4-yl)methanesulfonamide

-120-



The title compound was prepared as a white solid by the method of Example 115 using the subtitle product of Example 115 step iii) (1.0g), (R)-(-)-2-amino-1-butanol (1.5ml) and
5 trifluoroacetic acid (2ml). Yield: 0.19g.


MS APCI(+ve) 415 $[M+H]^+$

^1H NMR $\delta_{(\text{DMSO})}$ 10.52 (1H, s), 7.36 (1H, t), 7.17 (1H, t), 7.01 (1H, t), 5.81 (1H, s), 4.65 (1H, s), 4.32 (2H, s), 3.92 (1H, s), 3.42 - 3.37 (1H, m), 3.31 (3H, s), 3.34 - 3.29 (1H, m), 2.23 (3H, s), 1.65 - 1.56 (1H, m), 1.41 - 1.32 (1H, m), 0.84 (3H, t).

10

Example 118

N-(2-[[2-Fluoro-3-(trifluoromethyl)benzyl]thio]-6-[[*(1R)*]-2-hydroxy-1-methylethyl]-amino}pyrimidin-4-yl)methanesulfonamide



15

The title compound was prepared as pale yellow crystals by the method of Example 115 using the subtitle product of step iii) (0.21g) and (R)-alaninol (0.3ml). Yield: 0.12g.

MS APCI(+ve) 455 $[M+H]^+$

20 ^1H NMR $\delta_{(\text{CDCl}_3)}$ 7.71 (1H, t), 7.49 (1H, t), 7.19 (1H, t), 5.95 (1H, s), 5.07 (1H, s), 4.35 (2H, s), 3.74 - 3.69 (1H, m), 3.59 - 3.54 (1H, m), 3.15 (3H, s), 1.20 (3H, d).

The intermediates for this compound were prepared as follows:

i) 2-[[2-Fluoro-3-(trifluoromethyl)benzyl]thio}pyrimidine-4,6-diol

-121-

The subtitle compound was prepared as a yellow solid by the method of Example 39 step i) using 2-fluoro-3-(trifluoromethyl)benzyl bromide (2.0g), 2-mercaptopyrimidine-4,6-diol (1.12g) and potassium hydroxide (0.44g). Yield: 2.23g.

MS APCI(+ve) 321 [M+H]⁺

5

ii) 4,6-Dichloro-2-[(2-fluoro-3-(trifluoromethyl)benzyl)thio]pyrimidine

The subtitle compound was prepared as white crystals by the method of Example 39 step ii) using the subtitle product of step i) (2.23g), phosphorus oxychloride (10.4ml) and *N,N*-dimethylaniline (2.0ml). Yield: 1.7g.

10 ¹H NMR δ_(CDCl₃) 7.75 (1H, t), 7.52 (1H, t), 7.18 (1H, t), 7.06 (1H, s), 4.43 (2H, s).

iii) *N*-(6-Chloro-2-[(2-fluoro-3-(trifluoromethyl)benzyl)thio]pyrimidin-4-yl)-*N*-{2-(trimethylsilyl)ethoxy)methyl}methanesulfonamide


The subtitle compound was prepared as a colourless oil by the method of Example 39 step iii)

15 using the subtitle product of step ii) (0.57g), methanesulfonamide (0.15g), 60% sodium hydride (0.26g) and 2-(trimethylsilyl)ethoxymethyl chloride (0.3ml). Yield: 0.21g.

¹H NMR δ_(CDCl₃) 7.77 (1H, t), 7.53 (1H, t), 7.19 (1H, t), 7.12 (1H, s), 5.32 (2H, s), 4.45 (2H, s), 3.66 (2H, t), 3.32 (3H, s), 0.93 (2H, t), 0.00 (9H, s).

20 **Example 119**

***N*-(2-[(2,3-Difluoro-4-methylbenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)methanesulfonamide**



25

The title compound was prepared from a solution of the product of Example 43 step iv) (4ml) and quenching 2,3-difluoro-4-methylbenzyl bromide (1.46g) using the method described for Example 43 to give the title compound as a white solid. Yield: 4mg.


MS APCI(+ve) 419 [M+H]⁺

-122-

^1H NMR δ (CD_3OD) 7.27 - 7.23 (1H, m), 6.93 - 6.89 (1H, m), 5.56 (1H, s), 4.41 (2H, s), 3.91 - 3.81 (1H, m), 3.55 - 3.50 (1H, m), 3.48 - 3.43 (1H, m), 3.03 (3H, s), 2.25 (3H, s), 1.16 (3H, d).

5 **Example 120**

***N*-(2-[(2-Fluoro-3-methoxybenzyl)thio]-6-[(1*R*)-2-hydroxy-1-methylethyl]amino)-pyrimidin-4-yl)methanesulfonamide**



10

The title compound was prepared as a white foam by the method of Example 40 using the subtitle product of step vii) (0.24g) and trifluoroacetic acid (1.0ml). Yield: 0.11g.

MS APCI(+ve) 417 $[\text{M}+\text{H}]^+$

^1H NMR δ (DMSO) 7.24 (1H, bs), 7.06 (3H, m), 5.77 (1H, s), 4.70 (1H, bt), 4.33 (2H, t), 4.00
15 (1H, bs), 3.82 (3H, s), 3.40 (1H, m), 3.28 (1H, m), 3.20 (3H, s), 1.06 (3H, d).

The intermediates for this compound were prepared as follows:

i) 2-Fluoro-3-methoxybenzoic acid

Pentamethylenediethylenetetramine (31.2ml) was added to a solution of 2-fluoroanisole
20 (15.0g) in THF (450ml). The reaction mixture was cooled to -78°C and *n*-butyllithium
(59.6ml, 2.5M solution in hexanes) was added dropwise. Stirring was maintained for 2h
before the solution was added in a dropwise fashion to a flask containing solid carbon dioxide
pellets. Upon complete addition (30min) the mixture was allowed to warm to room
temperature before removal of the volatiles *in vacuo*. The residue was dissolved in 10%
25 sodium hydroxide solution (300ml) and extracted with Et_2O (3x). The aqueous was acidified
to pH 1 with concentrated hydrochloric acid before extracting with DCM. The organics were
washed with H_2O , dried (MgSO_4) and concentrated *in vacuo* to afford the subtitle compound
as a yellow solid. Yield: 7.1g.

^1H NMR δ (CDCl_3) 7.50 (1H, m), 7.12 (2H, m), 3.91 (3H, s).

ii) (2-Fluoro-3-methoxyphenyl)methanol

Lithium aluminium hydride (83.5ml, 1M solution in THF) was added dropwise to a suspension of the subtitle product of step i) (7.1g) in Et₂O (180ml) at a rate that maintained gentle reflux. Upon complete addition the reaction was stirred for 1.5h. 15% sodium hydroxide solution was added dropwise until no effervescence was observed. The resulting white precipitate was filtered and the filtrate diluted with H₂O (100ml). The organics were removed *in vacuo* and the residue extracted with Et₂O (100ml). The organics were washed with 2M sodium hydroxide solution (150ml), H₂O (150ml), brine (150ml), dried (MgSO₄) and concentrated *in vacuo* to afford the subtitle compound as a white crystalline solid. Yield: 5.5g.

¹H NMR δ (CDCl₃) 7.00 (3H, m), 4.77 (2H, d), 3.89 (3H, s), 1.77 (1H, t).

iii) 1-(Bromomethyl)-2-fluoro-3-methoxybenzene

Triphenylphosphine (11.1g) was added to a solution of the subtitle product of step ii) (5.0g) in DCM (200ml) followed by the portionwise addition of carbon tetrabromide (14.0g). The reaction was stirred for 4h before the addition of further triphenylphosphine (2.0g) and carbon tetrabromide (2.0g) and stirring maintained for 1h. The mixture was concentrated to 30ml volume and diluted in Et₂O (300ml). The solid precipitate was filtered and washed with Et₂O (3x) and the filtrate concentrated *in vacuo*. The residue was purified by column chromatography (10% Et₂O / iso-hexane) to afford the subtitle compound as a clear oil. Yield: 5.2g.

¹H NMR δ (CDCl₃) 7.05 (1H, m), 6.93 (2H, m), 4.52 (2H, s), 3.89 (3H, s).

iv) 2-[(2-Fluoro-3-methoxybenzyl)thio]pyrimidine-4,6-diol

The subtitle compound was prepared as a white solid by the method of Example 39 step i) using 1-(bromomethyl)-2-fluoro-3-methoxybenzene (4.5g), 2-mercaptopyrimidine-4,6-diol (2.96g) and potassium hydroxide (1.15g). Yield: 5.0g.

¹H NMR δ (DMSO) 7.10 (3H, m), 5.21 (1H, bs), 4.38 (2H, s), 3.83 (3H, s).

v) 4,6-Dichloro-2-[(2-fluoro-3-methoxybenzyl)thio]pyrimidine

-124-

The subtitle compound was prepared as a white solid by the method of Example 39 step ii) using the subtitle product of step iv) (4.91g), phosphorus oxychloride (42.6ml) and *N,N*-dimethylaniline (4.9ml). Yield: 4.1g.

$^1\text{H NMR } \delta_{(\text{DMSO})}$ 7.74 (1H, s), 7.09 (3H, m), 4.43 (2H, s), 3.83 (3H, s).

5

vi) *N*-{6-Chloro-2-[(2-fluoro-3-methoxybenzyl)thio]pyrimidin-4-yl}-*N*-{[2-(trimethylsilyl)ethoxy]methyl}methanesulfonamide

The subtitle compound was prepared as a colourless oil by the method of Example 39 step iii) using the subtitle product of step v) (2.0g), methanesulfonamide (0.60g), 60% sodium hydride (0.50g) and 2-(trimethylsilyl)ethoxymethyl chloride (1.11ml). Yield: 2.42g.

10

MS APCI(+ve) 509 $[\text{M}+\text{H}]^+$

vii) *N*-{6-Chloro-2-[(2-fluoro-3-methoxybenzyl)thio]pyrimidin-4-yl}-*N*-{[2-(trimethylsilyl)ethoxy]methyl}methanesulfonamide


15 A solution of the subtitle product of step vi) (0.3g) in (*R*)-alaninol (1.5ml) was stirred at 90°C for 2.5h. The reaction mixture was diluted in EtOAc (50ml) and washed with water, brine, dried (MgSO_4) and concentrated *in vacuo*. The residue was purified by column chromatography (50% EtOAc / iso-hexane) to afford the subtitle compound as a gum. Yield: 0.24g.

20 MS APCI(+ve) 547 $[\text{M}+\text{H}]^+$

Example 121

***N*-(2-[(2-Fluoro-3-methoxybenzyl)thio]-6-[[*(1R)*-1-(hydroxymethyl)propyl]amino]-pyrimidin-4-yl)methanesulfonamide**

25



-125-

The title compound was prepared as a white foam by the method of Example 40 using the subtitle product of step i) (0.24g) and trifluoroacetic acid (1.0ml). Yield: 0.11g.

MS APCI(+ve) 431 $[M+H]^+$

^1H NMR δ (DMSO) 7.10 (1H, bs), 7.06 (3H, m), 5.80 (1H, bs), 4.64 (1H, bs), 4.30 (2H, t), 3.85
5 (1H, bs), 3.82 (3H, s), 3.39 (1H, bm), 3.20 (3H, bs), 1.61 (1H, p), 1.36 (1H, p), 0.83 (3H, t).

The intermediates for this compound were prepared as follows:

i) *N*-(2-[(2-Fluoro-3-methoxy-benzyl)thio]-6-{[(1*R*)-1-(hydroxymethyl)propyl]amino}-pyrimidin-4-yl)-*N*-{2-(trimethylsilyl)ethoxy)methyl}methanesulfonamide


10 A solution of the subtitle compound of Example 120 step vi) (0.3g) in (2*R*)-2-aminobutan-1-ol (1.5ml) was stirred at 90°C for 2.5h. The reaction mixture was diluted in EtOAc (50ml) and washed with H₂O and brine, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography (50% EtOAc / iso-hexane) to afford the subtitle compound as a gum. Yield: 0.26g.

15 MS APCI(+ve) 561 $[M+H]^+$

Example 122

N-{6-{[(1*R*)-2-Hydroxy-1-methylethyl]amino}-2-[(3-methoxy-2-methylbenzyl)thio]-pyrimidin-4-yl}-1,2-dimethyl-1*H*-imidazole-4-sulfonamide

20



A solution of the subtitle compound of step i) (0.4g) and (*R*)-alaninol (0.11ml) in NMP (1.0ml) was stirred at 90°C for 2h. Upon cooling the reaction mixture was diluted with
25 acetonitrile (4ml) and 2M hydrochloric acid (1ml) and stirring maintained for 10min. The solvent was partially evaporated under reduced pressure and the reaction mixture diluted in EtOAc (50ml) and washed with H₂O (5ml), dried (MgSO₄) and concentrated *in vacuo* to afford a yellow oil. The residue was purified by reverse phase HPLC (gradient 90% to 5%

-126-

0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid.

Yield: 12mg.

MS APCI(+ve) 497 [M+H]⁺

¹H NMR δ (DMSO) 7.70 (1H, bs), 7.03 (4H, m), 5.88 (1H, bs), 4.69 (2H, bs), 3.90 (1H, bs), 3.54
5 (3H, s), 3.37 (1H, m), 3.28 (3H, s), 3.25 (1H, m), 2.26 (3H, s), 1.03 (3H, d).

The intermediates for this compound were prepared as follows:

i) *N*-{6-Chloro-2-[(3-methoxy-2-methylbenzyl)thio]pyrimidin-4-yl}-1,2-dimethyl-*N*-{[2-(trimethylsilyl)ethoxy]methyl}-1*H*-imidazole-4-sulfonamide


10 The subtitle compound was prepared as a colourless oil by the method of Example 39 step iii) using the subtitle product of Example 120 step v) (2.3g), 1,2-dimethyl-1*H*-imidazole-4-sulfonamide (0.60g), 60% sodium hydride (0.87g) and 2-(trimethylsilyl)ethoxymethyl chloride (1.5ml). Yield: 2.2g.

MS APCI(+ve) 646 [M+H]⁺

15

Example 123

N-{6-[(2-Hydroxy-1,1-dimethylethyl)amino]-2-[(3-methoxy-2-methylbenzyl)thio]pyrimidin-4-yl}-1,2-dimethyl-1*H*-imidazole-4-sulfonamide



20

The title compound was prepared as a white solid by the method of Example 122 using the subtitle product of Example 122 step i) (0.40g) and 2-amino-2-methylpropan-1-ol (0.20ml) then 2M hydrochloric acid (1.0ml). Yield: 13mg.


25 MS APCI(+ve) 511 [M+H]⁺

¹H NMR δ (DMSO) 7.28 (1H, bs), 7.04 (3H, m), 5.23 (1H, bs), 5.72 (1H, s), 4.23 (2H, s), 3.82 (3H, s), 3.52 (3H, s), 3.35 (2H, bs), 2.25 (3H, s), 1.18 (6H, s).

-127-

Example 124

***N*-{6-[[*(1R)*-1-(Hydroxymethyl)propyl]amino}-2-[(3-methoxy-2-methylbenzyl)thio]pyrimidin-4-yl}-1,2-dimethyl-1*H*-imidazole-4-sulfonamide**



5


The title compound was prepared as a white solid by the method of Example 122 using the subtitle product of Example 122 step i) (0.40g), and (*R*)-2-aminobutanol (0.19ml) then 2M hydrochloric acid (1.0ml). Yield: 46mg.

10 MS APCI(+ve) 511 [M+H]⁺

¹H NMR δ (DMSO) 7.72 (1H, bs), 7.03 (1H, bs), 7.00 (3H, bs), 5.92 (1H, bs), 4.62 (1H, bs), 4.26 (2H, s), 3.82 (4H, bs + s), 3.54 (3H, s), 3.36 (1H, m), 2.27 (3H, s), 1.57 (1H, m), 1.35 (1H, m), 0.81 (3H, t).

15 **Example 125**

***N*-(2-(Benzylthio)-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)-1,1,1-trifluoromethanesulfonamide**



20

Triflic anhydride (0.38ml) was added dropwise to a solution of the subtitle product of Example 3 step ii) (0.4g) and *N,N*-diisopropylethylamine (1.7ml) in DCM at -10°C. After 15min saturated sodium bicarbonate (10ml) was added and the organics recovered through extraction with DCM (2x10ml). The organics were combined, washed with H₂O, brine, dried

-128-

(MgSO₄) and concentrated *in vacuo*. The residue was dissolved in THF (10ml) and treated with tetrabutylammonium fluoride (5ml, 1M in THF) for 15mins before acidifying to pH 1 with 1M hydrochloric acid. EtOAc (10ml) was added before the organics were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column

5 chromatography (2% methanol / DCM) to afford a gum which was freeze dried from dioxane (20ml) to yield title compound as a foam. Yield: 0.37g.


MS APCI(+ve) 422 [M+H]⁺

¹H NMR δ_(DMSO) 13.30 (1H, s), 8.44 (1H, d), 7.42 (2H, d), 7.37 - 7.24 (3H, m), 6.20 (1H, s), 4.48 - 4.41 (2H, m), 4.28 - 4.16 (1H, m), 3.46 - 3.28 (2H, m), 1.09 (3H, d).

10

Example 126

N-(2-(Benzylthio)-5-chloro-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)methanesulfonamide



15

The title compound of Example 3 (0.4g) was dissolved in DCM (20ml) and treated with *N*-chlorosuccinimide (0.14g) for 2h. The volatiles were removed *in vacuo* and the residue purified by reverse phase HPLC ((gradient 95% to 20% 0.02M ammonium hydroxide /
20 acetonitrile) to yield the title compound as a white solid. Yield: 0.25g.


MS APCI(-ve) 402 [M-H]⁻

¹H NMR δ_(DMSO) 10.33 (1H, s), 7.41 (2H, d), 7.35 - 7.20 (3H, m), 6.76 (1H, d), 4.79 (1H, t), 4.39 - 4.29 (2H, m), 4.28 - 4.16 (1H, m), 3.50 - 3.31 (2H, m), 1.12 (3H, d).

25 Example 127

N-(2-(Benzylthio)-6-{[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)-2-chlorobenzenesulfonamide

-129-




2-Chlorophenylmethanesulfonyl chloride (0.17g) was added to a solution of the subtitle product of Example 3 step ii) (66mg) in pyridine (2ml) and *N,N*-dimethylaminopyridine (24mg). The reaction mixture was stirred for 18h. The volatiles were removed under reduced pressure and the residue diluted in THF (5ml) and treated with 2M hydrochloric acid (5ml) for 5min. The solvent was evaporated and the residue partitioned between DCM and treated with saturated sodium bicarbonate to pH neutral. The organic layer was washed with H₂O and brine. The organics were dried (MgSO₄) and concentrated to yield a solid. This material was purified by column chromatography (20% EtOAc / DCM) to yield the title compound as an orange solid. Yield: 19mg.

MS APCI(+ve) 464 [M+H]⁺

¹H NMR δ (DMSO) 10.88 (1H, s), 8.07 (1H, d), 7.66 (2H, m), 7.56 (1H, m), 7.27 (5H, m), 6.45 (1H, bd), 6.09 (1H, s), 5.91 (1H, s), 4.24 (2H, q), 3.85 (1H, bt), 3.30 (2H, m), 1.02 (3H, d).

Example 128

N-(2-[(3,4-Difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)methanesulfonamide



The title compound was prepared by the method of Example 27 using the subtitle product of step iii) (0.3g) and methanesulfonyl chloride (0.16ml). Yield: 77mg.

MS APCI(+ve) 405 [M+H]⁺

-130-

^1H NMR $\delta_{(\text{DMSO})}$ 7.51 (1H, m), 7.33 (2H, m), 5.78 (1H, s), 4.70 (1H, bs), 4.30 (2H, q), 3.95 (1H, bs), 3.33 (2H, m), 3.20 (3H, s), 1.05 (3H, d).

The intermediates for the above compound were prepared as follows:

5 **i) 6-Amino-2-[(3,4-difluorobenzyl)thio]pyrimidin-4-ol**

The subtitle compound was prepared according to the procedure of Example 1 step i) treating 4-amino-6-hydroxy-2-mercaptopyrimidine monohydrate (2.0g) with 3,4- difluorobenzyl bromide (2.66g) to afford the subtitle compound as a white solid. Yield: 3.35g.

^1H NMR $\delta_{(\text{DMSO})}$ 7.54 (1H, m), 7.32 (2H, m), 6.58 (2H, bs), 4.96 (1H, bs), 4.29 (2H, s).

10

ii) 6-Chloro-2-[(3,4-difluorobenzyl)thio]pyrimidin-4-amine

The subtitle compound was prepared from the product of step i) (3.35g) according to the procedure of Example 1 step ii) to afford the subtitle product as a green foam which was used directly in the subsequent step.

15 MS: APCI(+ve) 368 $[\text{M}+\text{H}]^+$


iii) *N*-((1*R*)-2-[[*tert*-Butyl(dimethyl)silyl]oxy]-1-methylethyl)-2-[(3,4-difluorobenzyl)-thio]pyrimidine-4,6-diamine

N,N-Diisopropylethylamine (4.9ml) was added to a solution of (*R*)-alaninol (5.0ml) and the
20 subtitle product of step ii) and stirred at 120°C for 7 days before partitioning between H₂O and DCM. The organics were washed with H₂O, brine, dried (MgSO₄) and concentrated *in vacuo* to afford a residue which was purified by column chromatography (8:1 EtOAc / iso-hexane). The residue was treated with imidazole (0.29g) and a solution of *tert*-butyldimethylsilyl chloride (0.63g) in DMF (1.5ml) and stirring maintained for 18h. The
25 reaction mixture was partitioned between EtOAc and H₂O and the organics recovered, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography (6:4 Et₂O / iso-hexane) to afford the subtitle compound as an orange gum. Yield: 0.61g.
MS: APCI(+ve) 441 $[\text{M}+\text{H}]^+$

30 **Example 129**

***N*-(2-(Benzylthio)-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonamide**

-131-




The title compound was prepared by the method of Example 127 using the subtitle product of Example 3 step ii) (0.2g) and 2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-sulfonyl chloride (0.49g). Yield: 84mg.

MS APCI(+ve) 486 [M+H]⁺

¹H NMR δ (DMSO-D₂O) 7.59 (1H, d), 7.55 (1H, s), 7.27 (6H, m), 5.65 (1H, s), 4.17 (2H, t), 4.01 (2H, s), 3.81 (1H, bs), 3.37 (1H, m), 3.24 (1H, m), 3.09 (2H, t), 2.84 (2H, t), 1.03 (3H, d).

10 Example 130

5-[[2-(Benzylthio)-6-[[1(1R)-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl]amino]sulfonyl]-2-furoic acid



15

Hydrogen chloride (2ml, 4M in dioxan) was added to the subtitle product of step ii) (30mg) and stirred for 2h. The volatiles were removed *in vacuo* and the residue purified by reverse phase HPLC (gradient 95% to 50% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 20mg.

20 MS APCI(+ve) 436 [M+H]⁺

¹H NMR δ (DMSO) 7.40 (2H, d), 7.30 (3H, m), 6.97 (1H, d), 6.91 (1H, d), 5.85 (1H, s), 4.34 (2H, q), 4.02 (1H, bs), 3.39 (1H, m), 3.30 (1H, m), 1.07 (3H, d).

The intermediates for this compound were prepared as follows:

-132-

i) Methyl 5-[(2-(benzylthio)-6-(((1*R*)-2-[[*tert*-butyl(dimethyl)silyl]oxy]-1-methylethyl)amino)pyrimidin-4-yl)amino)sulfonyl]-2-furoate

Methyl 5-(chlorosulfonyl)-2-furoate (0.54g) was added to the solution of the subtitle product of Example 3 step ii) (0.5g) in pyridine (15ml) and *N,N*-dimethylaminopyridine (0.15g). The reaction mixture was stirred for 18h. A further aliquot of the sulfonyl chloride (0.27g) was added and stirring maintained for a further 18h. The solvent was evaporated and the residue purified by column chromatography (25% EtOAc / iso-hexane) to yield the subtitle compound as a yellow glass. Yield: 0.24g.

¹H NMR δ (DMSO) 7.38 (7H, m), 7.09 (1H, bs), 5.95 (1H, s), 4.42 (2H, s), 4.26 (1H, bs), 3.87 (3H, s), 3.54 (2H, bm), 1.12 (3H, d), 0.85 (9H, s), 0.01 (6H, m).


ii) 5-[(2-(Benzylthio)-6-(((1*R*)-2-[[*tert*-butyl(dimethyl)silyl]oxy]-1-methylethyl)amino)pyrimidin-4-yl)amino)sulfonyl]-2-furoic acid

Lithium hydroxide (33mg) was added to a solution of the subtitle product of step i) (0.23g) in THF / H₂O (1ml / 1ml) and stirring maintained for 1h. The THF was removed *in vacuo* and the residue neutralised with AcOH before extracting with EtOAc. The organics were then washed with H₂O, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by reverse phase HPLC (gradient 95% to 20% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as an off-white solid. Yield: 0.14g.

¹H NMR δ (DMSO) 7.13 (6H, m), 6.84 (2H, d), 5.72 (1H, s), 4.26 (2H, s), 3.93 (1H, bs), 3.58 (1H, m), 3.34 (1H, m), 1.07 (3H, d), 0.84 (9H, s), 0.00 (6H, s).

Example 131

***N*-(2-(Benzylthio)-6-(((1*R*)-2-hydroxy-1-methylethyl)amino)pyrimidin-4-yl)-5-(piperazin-1-ylcarbonyl)furan-2-sulfonamide**



-133-

Hydrogen chloride (2ml, 4M in dioxan) was added to the subtitle product of step i) (0.12g) and stirred for 2h. The volatiles were removed *in vacuo* and the residue purified by reverse phase HPLC (gradient 95% to 50% 0.02M ammonium hydroxide / acetonitrile) to yield the
5 title compound as a white solid. Yield: 68mg.

MS APCI(+ve) 533 [M+H]⁺

¹H NMR δ (DMSO) 7.35 (2H, d), 7.24 (3H, m), 7.02 (1H, d), 6.86 (1H, d), 5.68 (1H, s), 4.21 (2H, t), 3.84 (1H, bs), 3.73 (4H, bs), 3.38 (1H, m), 3.25 (1H, m), 3.01 (4H, bs), 1.05 (3H, d).

10 The intermediates for this compound were prepared as follows:

i) *tert*-Butyl 4-[(2-(benzylthio)-6-[(1*R*)-2-[[*tert*-butyl(dimethyl)silyl]oxy]-1-methyl-ethyl)amino]-pyrimidin-4-yl]amino)sulfonyl]piperazine-1-carboxylate

tert-Butylpiperazine-1-carboxylate (45mg), *N*-hydroxybenzotriazole (33mg), and then dicyclohexylcarbodiimide (50mg) were added to a solution of the subtitle product of Example


15 130 step ii) (0.14g) in DCM (5ml). After 1h the reaction was filtered and washed well with DCM. The combined filtrates were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography (40% EtOAc / iso-hexane) to yield the subtitle compound as a white solid. Yield: 0.17g.

MS APCI(+ve) 747 [M+H]⁺

20

Example 132

5-[(2-(Benzylthio)-6-[(1*R*)-2-hydroxy-1-methylethyl]amino}pyrimidin-4-yl)amino)sulfonyl]-*N,N*-dimethyl-2-furamide



2M Hydrochloric acid (10ml) was added to a solution of the subtitle product of step i) (0.20g) in THF (10ml) and stirred for 3h. The volatiles were removed *in vacuo* and the residue was

-134-

extracted with DCM. The organics were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by reverse phase HPLC (gradient 95% to 50% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a white solid. Yield: 31mg.

5 MS APCI(-ve) 490 [M-H]⁻

¹H NMR δ (DMSO) 7.39 (2H, d), 7.28 (3H, m), 7.05 (2H, bs), 5.93 (1H, vbs), 4.78 (1H, bs), 4.35 (2H, bs), 4.13 (1H, vbs), 3.40 (2H, m), 3.09 (3H, bs), 2.95 (3H, bs), 1.07 (3H, d).

The intermediates for this compound were prepared as follows:

10 i) 5-[(2-(Benzylthio)-6-(((1*R*)-2-[[*tert*-butyl(dimethyl)silyl]oxy}-1-methylethyl)amino)-pyrimidin-4-yl)amino)sulfonyl]-*N,N*-dimethyl-2-furamide


A solution of the subtitle product of Example 130 step i) (0.37g) in 40% aqueous dimethylamine (4.2ml) was stirred for 18h. The volatiles were removed *in vacuo* and the residue extracted with EtOAc. The organics were washed with brine, dried (MgSO₄) and

15 concentrated *in vacuo*. The residue purified by column chromatography (50% EtOAc / iso-hexane) to yield the subtitle compound as a yellow gum. Yield: 0.14g.

MS APCI(+ve) 606 [M+H]⁺

Example 133

20 *N*-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-(((1*R*)-2-hydroxy-1-methylethyl)amino)pyrimidin-4-yl)-*cis*-3,5-dimethylpiperazine-1-sulfonamide



25 The title compound was prepared as a white solid by the method of Example 39 using the subtitle product of step ii) (0.37g), and (*R*)-alaninol (1ml). Yield: 6mg.

MS APCI(+ve) 519 [M+H]⁺

-135-

¹H NMR δ (DMSO) 7.59 (1H, t), 7.47 (1H, p), 7.15 (1H, s), 5.80 (1H, s), 4.67 (1H, t), 4.35 (2H, s), 3.90 (1H, bs), 3.57 (2H, s), 3.38 (4H, m), 2.24 (2H, t), 1.04 (3H, d), 0.94 (6H, d).

The intermediates for this compound were prepared as follows:

5 i) ***cis*-3,5-dimethylpiperazine-1-sulfonamide**

A solution of *cis*-2,6-dimethylpiperazine (5.0g) and sulfamide (10.0g) in 1,4-dioxane (100ml) was stirred for 72h at 110°C. The volatiles were removed *in vacuo* and the residue suspended in EtOAc. The filtrate was evaporated to a yellow solid (4.3g). 1g of this material was dissolved in methanol and applied to an SCX cartridge (10g). The cartridge was washed with

- 10 50% aqueous methanol (200ml) before the subtitle product was eluted with 5% ammonium hydroxide solution / methanol (200ml). The solvent was removed under reduced pressure to yield the subtitle compound as a yellow solid. Yield: 0.46g.

MS APCI(+ve) 194 [M+H]⁺

15 ii) ***N*-(6-chloro-2-[(3-chloro-2-fluorobenzyl)thio]-pyrimidin-4-yl)-*cis*-3,5-dimethylpiperazine-1-sulfonamide**

60 % Sodium hydride (0.19g) was added to a stirred solution of the subtitle product of step i) (0.45g) in DMF (4.2ml) at 0°C. The cooling bath was removed for 15min before recooling to 0°C and addition of a solution of the subtitle product of Example 31 step iii) (0.76g) in DMF

- 20 (2ml). After stirring at room temperature for 3h the mixture was acidified with 2M hydrochloric acid to pH 4 and the volatiles were removed *in vacuo*. The residue was dissolved in methanol and applied to an SCX cartridge (10g). The cartridge was washed with methanol (200ml) before the subtitle product was eluted with 10% triethylamine / methanol (300ml). The solvent was removed under reduced pressure and the residue triturated from
- 25 Et₂O to yield the subtitle compound as a yellow solid. Yield: 0.37g.


MS APCI(+ve) 480/482/484 [M+H]⁺

Example 134

***N*-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[(1*R*)-2-hydroxy-1-**

- 30 **methylethyl]amino}pyrimidin-4-yl)-4-ethylpiperazine-1-sulfonamide**

-136-



A solution of the subtitle product of step ii) (1.92g) in (*R*)-alaninol (5ml) was heated at 80°C for 72h. The reaction mixture was then diluted in methanol and purified by reverse phase HPLC (gradient 95% to 60% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a yellow glass. Yield: 90mg.

MS APCI(+ve) 519 [M+H]⁺

¹H NMR δ (DMSO) 7.59 (1H, t), 7.47 (1H, t), 7.16 (2H, bt), 5.86 (1H, s), 4.69 (1H, bm), 4.36 (2H, t), 3.91 (1H, vbs), 3.38 (2H, m), 3.12 (4H, bs), 2.28 (6H, m), 1.05 (3H, d), 0.96 (3H, t).

10

The intermediates for this compound were prepared as follows:

i) 4-Ethylpiperazine-1-sulfonamide

A solution of *N*-ethylpiperazine (5.0g) and sulfamide (10.0g) in 1,4-dioxane (100ml) was stirred for 72h at 110°C. The volatiles were removed *in vacuo* and 5g of the residue was

15 dissolved in methanol and applied to an SCX cartridge (70g). The cartridge was washed with 50% aqueous methanol (200ml) before the subtitle product was eluted with 10% triethylamine / methanol (100ml). The solvent was removed under reduced pressure to yield the subtitle compound as a pale beige solid. Yield: 3.0g.

MS APCI(-ve) 192 [M-H]⁻

20

ii) *N*-{6-Chloro-2-[(3-chloro-2-fluorobenzyl)thio]pyrimidin-4-yl}-4-ethylpiperazine-1-sulfonamide

The subtitle compound was prepared as an orange gum by the method of Example 133 step ii) using the subtitle product of step i) (3.0g), 60 % sodium hydride (1.24g) and the subtitle


25 product of Example 31 step iii) (5.0g). Yield: 0.73g.

MS APCI(+ve) 480 [M+H]⁺

-137-

Example 135

***N*-{2-[(3-Chloro-2-fluorobenzyl)thio]-6-[(2-hydroxy-1,1-dimethylethyl)amino]-pyrimidin-4-yl}-*cis*-3,5-dimethylpiperazine-1-sulfonamide**



5

A solution of the subtitle product of step i) (0.26g) in 2-amino-2-methylpropanol (1ml) was heated at 90°C for 3.5h and then 55°C for 72h. The reaction mixture was then diluted in EtOAc and washed with H₂O, dried (Na₂SO₄) and concentrated *in vacuo*. The residue was
 10 diluted in trifluoroacetic acid (2ml) and stirred for 15min before removal of the volatiles *in vacuo* and azeotroping the residue with toluene (2x). The crude material and purified by reverse phase HPLC (gradient 95% to 50% 0.02M ammonium hydroxide / acetonitrile) to yield the title compound as a solid. Yield: 48mg.

MS APCI(+ve) 533 [M+H]⁺

15 ¹H NMR δ (DMSO) 7.58 (1H, t), 7.47 (1H, t), 7.17 (1H, t), 5.88 (1H, s), 4.36 (2H, s), 3.46 (4H, m), 2.73 (2H, bs), 2.30 (2H, t), 1.21 (6H, s), 0.97 (6H, d).

The intermediates for this compound were prepared as follows:

i) ***N*-{6-Chloro-2-[(3-chloro-2-fluorobenzyl)thio]pyrimidin-4-yl}-*cis*-3,5-dimethyl-*N*-{[2-(trimethylsilyl)ethoxy]methyl}piperazine-1-sulfonamide**
 20


The subtitle compound was prepared as a yellow gum by the method of Example 39 step iii) using the subtitle product of Example 133 step i) (2.6g), the subtitle product of Example 31 step iii) (4.35g), 60% sodium hydride (0.99g) and 2-(trimethylsilyl)ethoxymethyl chloride (2.38ml). Yield: 3.4g.

25 MS APCI(+ve) 610 [M+H]⁺

-138-

Example 136

***N*-{6-[[*(1R)*-2-Hydroxy-1-methylethyl]amino]-2-*(R,S)*-[(1-phenylethyl)thio]pyrimidin-4-yl}methanesulfonamide**



5

Methanesulfonyl chloride (0.16ml) was added to a stirred solution of the subtitle product of step ii) (0.29g) and *N,N*-diisopropylethylamine (0.36ml) in DCM (5ml). After stirring for 18h the volatiles were removed under reduced pressure and the residue diluted in THF (8ml) and
 10 treated with 1M sodium hydroxide solution (4.2ml). After 6h 2M hydrochloric acid was added to pH 1 and stirring maintained for 3 days. The reaction mixture was then neutralised with saturated sodium bicarbonate solution and the product extracted with DCM. The organics were washed with H₂O, dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by reverse phase HPLC (gradient 95% to 25% 0.02M ammonium hydroxide /
 15 acetonitrile) to yield the title compound as a white solid. Yield: 90mg.

MS APCI(+ve) 383 [M+H]⁺

¹H NMR δ_(DMSO) 7.45 (2H, d), 7.33 (2H, t), 7.24 (1H, m), 5.77 (1H, bs), 4.93 (1H, q), 4.71 (1H, bs), 3.41 (1H, m), 3.30 (1H, m), 3.23 (3H, bs), 1.67 (3H, dd), 1.07 (3H, dd).

20 The intermediates for the title compound were prepared as follows:

i) 6-Amino-2-[(1-phenylethyl)thio]pyrimidin-4-ol

The subtitle compound was prepared according to the procedure of Example 1 step i) treating 4-amino-6-hydroxy-2-mercaptopyrimidine monohydrate (5.0g) with α -methylbenzyl bromide (5.74g) to afford the subtitle compound which was used directly in the subsequent step.

25 MS APCI(+ve) 352 [M+H]⁺

ii) *N*-((*1R*)-2-[[*tert*-butyl(dimethyl)silyl]oxy]-1-methylethyl)-2-*(R,S)*-[(1-phenylethyl)thio]-pyrimidine-4,6-diamine


-139-

The subtitle compound was prepared from the product of step i) according to the procedure of Example 1 step ii) to afford the subtitle product as a green foam which was then diluted in (*R*)-alaninol (12.2ml), *N,N*-diisopropylethylamine (11.8ml) and NMP (16ml) and stirred at 130°C for 3 days before partitioning between H₂O and DCM. The organics were washed with H₂O, dried (MgSO₄) and concentrated *in vacuo* to afford a residue which was purified by column chromatography (8:2 EtOAc / iso-hexane). The residue was treated with imidazole (2.7g) and a solution of *tert*-butyldimethylsilyl chloride (5.95g) in DMF (30ml) and stirring maintained for 18h. The reaction mixture was partitioned between EtOAc and H₂O and the organics recovered, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography (6:4 Et₂O / iso-hexane) to afford the subtitle compound as a gum. Yield: 1.3g.

MS: APCI(+ve) 419 [M+H]⁺

Example 137

N-{6-[[[(1*R*)-2-hydroxy-1-methylethyl]amino]-2-[(2,3,4-trifluorobenzyl)thio]-pyrimidin-4-yl]methanesulfonamide




The title product was prepared from a solution of the product of Example 43 step iv) (4ml) and quenching with 2,3,4-trifluorobenzyl bromide (0.5g) using the method described for Example 43 to give the title compound as a white foam. Yield: 32mg.

MS APCI(+ve) 423 [M+H]⁺

Example 138

N-[2-[[[(3-Chloro-2-fluorophenyl)methyl]thio]-6-[(*R*)-(2-hydroxy-1-methylethyl)amino]-4-pyrimidinyl]-1-methyl-1*H*-imidazole-4-sulfonamide

-140-



The title compound was prepared as a white solid by the method of Example 37 from the product of Example 27 step iii) (1.4g) using 1-methyl-1H-imidazole-4-sulfonyl chloride (1.0g). Purification was by column chromatography (DCM / methanol / AcOH 190:10:1).

5 Yield: 1.0g.

MS APCI(+ve) 488 [M+H]⁺


¹H NMR $\delta_{(\text{DMSO})}$ 7.85 (1H, bs), 7.75 (1H, s), 7.50 (2H, m), 7.18 (1H, m), 5.91 (1H, m), 4.36 (2H, s), 3.60 (3H, s), 3.30 (2H, m), 1.10 (3H, d).

-141-

CLAIMS

1. A compound of formula (1), pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof:

5



wherein R¹ is a group selected from C₃₋₇carbocyclyl, C₁₋₈alkyl, C₂₋₆alkenyl and C₂₋₆alkynyl;

- 10 wherein the group is optionally substituted by 1, 2 or 3 substituents independently selected from fluoro, nitrile, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹, phenyl or heteroaryl; wherein phenyl and heteroaryl are optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R⁹,
- 15 C₁₋₆alkyl and trifluoromethyl;

wherein R² is C₃₋₇carbocyclyl, optionally substituted by 1, 2 or 3 substituents independently selected from:

- (a) fluoro, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -SO₂R¹⁰, -SO₂NR⁵R⁶,
 20 -NR⁸SO₂R⁹;
- (b) a 3-8 membered ring optionally containing 1, 2 or 3 atoms selected from O, S, -NR⁸ and whereby the ring is optionally substituted by C₁₋₃alkyl or fluoro; or
- (c) phenyl or heteroaryl, each of which is optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, -OR⁴, -NR⁵R⁶, -CONR⁵R⁶, -NR⁸COR⁹, -
 25 SO₂NR⁵R⁶, -NR⁸SO₂R⁹, C₁₋₆alkyl and trifluoromethyl;

or R² is a group selected from C₁₋₈alkyl, C₂₋₆alkenyl or C₂₋₆alkynyl wherein the group is substituted by 1, 2 or 3 substituents independently selected from hydroxy, amino, C₁₋₆alkoxy,

-142-

C₁₋₆alkylamino, di(C₁₋₆alkyl)amino, *N*-(C₁₋₆alkyl)-*N*-(phenyl)amino, *N*-C₁₋₆alkylcarbamoyl, *N,N*-di(C₁₋₆alkyl)carbamoyl, *N*-(C₁₋₆alkyl)-*N*-(phenyl)carbamoyl, carboxy, phenoxycarbonyl, -NR⁸COR⁹, -SO₂R¹⁰, -SO₂NR⁵R⁶ and -NR⁸SO₂R⁹;

5 wherein R³ is hydrogen or independently R²;

R⁴ is hydrogen or a group selected from C₁₋₆alkyl and phenyl, wherein the group is optionally substituted by 1 or 2 substituents independently selected from halo, phenyl, -OR¹¹ and -NR¹²R¹³;

10

R⁵ and R⁶ are independently hydrogen or a group selected from C₁₋₆alkyl and phenyl wherein the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, phenyl, -OR¹⁴, -NR¹⁵R¹⁶, -COOR¹⁴, -CONR¹⁵R¹⁶, -NR¹⁵COR¹⁶, -SO₂R¹⁰, -SONR¹⁵R¹⁶ and NR¹⁵SO₂R¹⁶

15 or

R⁵ and R⁶ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocyclic ring system optionally containing a further heteroatom selected from oxygen and nitrogen atoms, which ring is optionally substituted by 1, 2 or 3 substituents independently selected from phenyl, -OR¹⁴, -COOR¹⁴, -NR¹⁵R¹⁶, -CONR¹⁵R¹⁶,
20 -NR¹⁵COR¹⁶, -SO₂R¹⁰, -SONR¹⁵R¹⁶, NR¹⁵SO₂R¹⁶ or C₁₋₆alkyl (optionally substituted by 1 or 2 substituents independently selected from halo, -NR¹⁵R¹⁶ and -OR¹⁷ groups);

R¹⁰ is hydrogen or a group selected from C₁₋₆alkyl or phenyl, wherein the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, phenyl, -OR¹⁷ and -

25 NR¹⁵R¹⁶; and

each of R⁷, R⁸, R⁹, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷ is independently hydrogen, C₁₋₆alkyl or phenyl;

30 X is hydrogen, halo, cyano, nitro, hydroxy, C₁₋₆alkoxy (optionally substituted by 1 or 2 substituents selected from halo, -OR¹¹ and -NR¹²R¹³), -NR⁵R⁶, -COOR⁷, -NR⁸COR⁹, thio, C₁₋₆alkylthio (optionally substituted by 1 or 2 substituents selected from halo, -OR¹⁷, -NR¹⁵R¹⁶), -SO₂R¹⁰ or a group selected from C₃₋₇carbocyclyl, C₁₋₈alkyl, C₂₋₆alkenyl or C₂₋₆alkynyl,

-143-

wherein the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$ and $-NR^8SO_2R^9$;

5 R^x is trifluoromethyl, $-NR^5R^6$, phenyl, naphthyl, monocyclic or bicyclic heteroaryl wherein a heteroring may be partially or fully saturated and one or more ring carbon atoms may form a carbonyl group, and wherein each phenyl or heteroaryl group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COR^7$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$, $-NR^8SO_2R^9$, C_{1-6} alkyl or

10 trifluoromethyl;

or R^x is a group selected from C_{3-7} carbocyclyl, C_{1-8} alkyl, C_{2-6} alkenyl and C_{2-6} alkynyl whereby the group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COR^7$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$, $-NR^8SO_2R^9$, phenyl or heteroaryl; and wherein each phenyl or heteroaryl group is optionally

15 substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COR^7$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$, $-NR^8SO_2R^9$, C_{1-6} alkyl or trifluoromethyl;

or R^x and X together form a 4 to 8-membered sulfonamide ring optionally substituted by 1, 2 or 3 substituents independently selected from halo, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COOR^7$, $-$

20 NR^8COR^9 , $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$, $-NR^8SO_2R^9$, phenyl or heteroaryl; wherein phenyl and heteroaryl are optionally substituted by 1, 2 or 3 substituents independently selected from halo, cyano, nitro, $-OR^4$, $-NR^5R^6$, $-CONR^5R^6$, $-COOR^7$, $-NR^8COR^9$, $-SR^{10}$, $-SO_2R^{10}$, $-SO_2NR^5R^6$, $-NR^8SO_2R^9$, C_{1-6} alkyl and trifluoromethyl.

25 2. A compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof according to claim 1 wherein R^2 is C_{1-8} alkyl optionally substituted by 1 or 2 hydroxy substituents.

3. A compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester
30 thereof according to claim 1 wherein R^1 is benzyl optionally substituted by 1, 2 or 3 substituents independently selected from fluoro, chloro, bromo, methoxy, methyl and trifluoromethyl.

-144-

4. A compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof wherein R³ is hydrogen;

5. A compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof wherein X is hydrogen

6. A compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof wherein R^x is methyl, 1-methylimidazolyl, 1,2-dimethylimidazolyl, *N,N*-dimethylamino, azetidiny, pyrrolidiny, morpholiny and piperidiny.

10

7. A compound selected from the group consisting of:

N-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)methanesulfonamide

N-[2-[(3-Chloro-2-fluorobenzyl)thio]-6-[(2-hydroxy-1-methylethyl)amino]-4-pyrimidinyl]-4-

15 morpholinesulfonamide

N-[2-[(3-Chloro-2-fluorophenyl)methyl]thio]-6-[(2-hydroxy-1-methylethyl)amino]-4-pyrimidinyl]-1,2-dimethyl-1*H*-imidazole-4-sulfonamide

N-(2-[(2,3-Difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)piperidine-1-sulfonamide

20 *N*-(2-[(2,3-Difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)pyrrolidine-1-sulfonamide

N-(2-[(2,3-Difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)azetidine-1-sulfonamide

N-{6-[[*(1R)*-2-Hydroxy-1-methylethyl]amino]-2-[(2,3,4-trifluorobenzyl)thio]-pyrimidin-4-yl}morpholine-4-sulfonamide

25

N-(2-[(2,3-Difluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]pyrimidin-4-yl)morpholine-4-sulfonamide

N-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)azetidine-1-sulfonamide

30 *N*-{6-[[*(1R)*-2-Hydroxy-1-methylethyl]amino]-2-[(2,3,4-trifluorobenzyl)thio]-pyrimidin-4-yl}azetidine-1-sulfonamide

N-(2-[(3-Chloro-2-fluorobenzyl)thio]-6-[[*(1R)*-2-hydroxy-1-methylethyl]amino]-pyrimidin-4-yl)-*N,N*-dimethylsulfamide

-145-

N-[2-[[[(3-Chloro-2-fluorophenyl)methyl]thio]-6-[(*R*)-(2-hydroxy-1-methylethyl)amino]-4-pyrimidinyl]-1-methyl-1*H*-imidazole-4-sulfonamide;

and a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof.


- 5 8. A compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof according to any one of claims 1 to 7 for use as a medicament.
9. A compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof according to any one of claims 1 to 7 for use as a medicament for the treatment of
- 10 asthma, allergic rhinitis, COPD, inflammatory bowel disease, osteoarthritis, osteoporosis, rheumatoid arthritis, or psoriasis..
10. A compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof according to any one of claims 1-7, for use as a medicament for the treatment of
- 15 cancer.
11. The use of a compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, according to any one of claims 1 to 7 in the manufacture of a medicament for the treatment of human diseases or conditions in which modulation of
- 20 chemokine receptor activity is beneficial.
12. The use of a compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, according to any one of claims 1 to 7 in the manufacture of a medicament for the treatment of asthma, allergic rhinitis, COPD, inflammatory bowel
- 25 disease, irritable bowel syndrome, osteoarthritis, osteoporosis, rheumatoid arthritis, or psoriasis.
13. The use of a compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, according to any one of claims 1 to 7 in the manufacture of a
- 30 medicament for the treatment of cancer.

-146-

14. A pharmaceutical composition comprising a compound, pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof according to any one of claims 1 to 7; and a pharmaceutically-acceptable diluent or carrier.

5 15. A process for the preparation of a compound according to claim 1 comprising the steps of:

a) treating a compound of formula (2):




10

wherein R^1 , R^2 , R^3 and X are as defined in claim 1, with sulfonyl chlorides (R^xSO_2Cl where R^x is as defined in claim 1;

or

b) treating a compound of formula (7):




15

wherein R^1 , R^x and X are as defined in formula (1), L is a halogen and Y is either hydrogen or a protecting group with nucleophilic amines of the type NR^2R^3 as defined in formula (1) in

20 the presence or absence of a suitable base and solvent;

or

c) treating a compound of formula (8):



-147-

wherein R^1 , R^x and X are as defined in formula (1) and L is halogen, with sulfonamides of formula $R^xSO_2NH_2$ where R^x is as defined in formula (1) except NR^5R^6 in the presence of a suitable base and solvent.

and

- 5 independently for each of process variants a), b) or c), optionally thereafter (i), (ii), (iii), (iv) or (v) in any order:
 - i) removing any protecting groups;
 - ii) converting the compound of formula (1) into a further compound of formula (1)
 - iii) forming a salt
 - 10 iv) forming a prodrug
 - v) forming an *in vivo* hydrolysable ester.
16. A combination therapy which comprises administering a compound of formula (1) or a pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, or a
- 15 pharmaceutical composition or formulation comprising a compound of formula (1), concurrently or sequentially with other therapy and/or another pharmaceutical agent.
17. A combination therapy as claimed in claim 16 for the treatment of asthma, allergic rhinitis, COPD, inflammatory bowel disease, irritable bowel syndrome, osteoarthritis,
- 20 osteoporosis, rheumatoid arthritis, or psoriasis.
18. A combination therapy as claimed in claim 16 for the treatment of cancer.
19. A pharmaceutical composition which comprises a compound of formula (1) or a
- 25 pharmaceutically acceptable salt, solvate or *in vivo* hydrolysable ester thereof, in conjunction with another pharmaceutical agent.
20. A pharmaceutical composition as claimed in claim 19 for the treatment of asthma, allergic rhinitis, COPD, inflammatory bowel disease, irritable bowel syndrome, osteoarthritis,
- 30 osteoporosis, rheumatoid arthritis, or psoriasis.
21. A pharmaceutical composition as claimed in claim 19 for the treatment of cancer.

INTERNATIONAL SEARCH REPORT

Application No
PCT/GB 03/03175

A. CLASSIFICATION OF SUBJECT MATTER					
IPC 7	C07D239/48	C07D239/69	C07D401/12	C07D403/12	C07D405/12
	C07D409/12	C07D413/12	C07D413/14	C07D417/12	A61K31/5377
	A61K31/506	A61P11/00	A61P29/00	A61P35/00	
According to International Patent Classification (IPC) or to both national classification and IPC					
B. FIELDS SEARCHED					
Minimum documentation searched (classification system followed by classification symbols)					
IPC 7	C07D	A61K			
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched					
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)					
EPO-Internal, WPI Data, PAJ, CHEM ABS Data					
C. DOCUMENTS CONSIDERED TO BE RELEVANT					
Category *	Citation of document, with indication, where appropriate, of the relevant passages				Relevant to claim No.
X	GB 1 042 295 A (CHUGAI PHARMACEUTICAL CO LTD) 14 September 1966 (1966-09-14) page 2, column 2, line 35 - line 37; claims 1,11; example 20 ---				1,2,5,8
X	WO 01 25242 A (HUNT SIMON FRASER ;WALTERS IAIN ALISTAIR STEWART (GB); ASTRAZENECA) 12 April 2001 (2001-04-12) page 12, line 28 -page 14, line 22; claims 1-6,10-19; examples 6,10 ---				1-21
X	WO 00 09511 A (ASTRA PHARMA PROD ;KINCHIN ELIZABETH (GB); AUSTIN RUPERT (GB); BAX) 24 February 2000 (2000-02-24) page 30, line 3 -page 32, line 4; claims 1-7,11-17; examples 231-235,272,294,295 ---				1-21
	-/--				
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.					
* Special categories of cited documents : *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *Z* document member of the same patent family					
Date of the actual completion of the international search			Date of mailing of the international search report		
16 October 2003			04/11/2003		
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016			Authorized officer Hanisch, I		

INTERNATIONAL SEARCH REPORT

Application No
PCT/GB 03/03175

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 01 58906 A (BONNERT ROGER ;HUNT FRASER (GB); WILLIS PAUL (GB); ASTRAZENECA AB) 16 August 2001 (2001-08-16) page 12, line 21 -page 15, line 12; claims 1-7,9-17; examples</p> <p>-----</p>	1-21

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 03/03175

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: —
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.2

It is noted that the application refers to "prodrugs" and "in vivo hydrolysable esters". Both are functional definitions which attempt to define a chemical compound in terms of a result to be achieved. This is not allowable (Article 6 PCT). The said terms have not been searched and should be deleted. "Prodrugs" and "in vivo hydrolysable esters" are functional definitions without a specific technical guidance for the selection of the suitable derivatives in the description and without proven general knowledge to show which derivatives are suitable prodrugs or which esters are indeed hydrolysable under in vivo conditions. The term could be seen as a mere invitation to the skilled person to perform a research program in order to find the suitable variants. Page 5, line 33-page 6, line 2 confirms this by stating "such esters can be identified by administering intravenously to a test animal the compound under test and subsequently examining the test animal's body fluid". In such a situation, when the invention cannot be carried out over the whole claimed area without imposing an undue burden, the disclosure may be considered insufficient in the sense of Article 5 PCT, even when simple in vivo or in vitro tests are available to determine whether or not a particular compound is covered by the claims.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Application No

PCT/GB 03/03175

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 1042295	A	14-09-1966	BE 641881 A	16-04-1964
			CH 460022 A	31-07-1968
			DE 1445609 A1	09-01-1969
			FR 3106 M	
			FR 1383287 A	24-12-1964
			NL 127990 C	
			NL 302745 A	
			US 3328395 A	27-06-1967
WO 0125242	A	12-04-2001	AU 7304900 A	10-05-2001
			BR 0014334 A	11-06-2002
			CA 2385269 A1	12-04-2001
			CN 1402730 T	12-03-2003
			CZ 20021113 A3	12-06-2002
			EE 200200174 A	15-04-2003
			EP 1222195 A1	17-07-2002
			EP 1348709 A2	01-10-2003
			WO 0125242 A1	12-04-2001
			HU 0204246 A2	28-04-2003
			JP 2003511384 T	25-03-2003
			NO 20021448 A	22-05-2002
			NZ 517880 A	26-09-2003
WO 0009511	A	24-02-2000	AT 231872 T	15-02-2003
			AU 5662599 A	06-03-2000
			CA 2338600 A1	24-02-2000
			DE 69905170 D1	06-03-2003
			DK 1104425 T3	23-06-2003
			EP 1104425 A1	06-06-2001
			ES 2190246 T3	16-07-2003
			JP 2002522544 T	23-07-2002
			NZ 509457 A	26-07-2002
			WO 0009511 A1	24-02-2000
WO 0158906	A	16-08-2001	GB 2359079 A	15-08-2001
			GB 2359080 A	15-08-2001
			AU 3255601 A	20-08-2001
			EP 1257555 A1	20-11-2002
			JP 2003522191 T	22-07-2003
			WO 0158906 A1	16-08-2001